



HAL
open science

Clinical and Molecular Spectrum Associated with COL6A3 c.7447A>G p.(Lys2483Glu) Variant: Elucidating its Role in Collagen VI-related Myopathies

Rocío Villar-Quiles, Sandra Donkevoort, Alix de Becdelievre, Corine Gartioux, Valérie Jobic, A. Reghan Foley, Riley Mccarty, Ying Hu, Rita Menassa, Laurence Michel, et al.

► To cite this version:

Rocío Villar-Quiles, Sandra Donkevoort, Alix de Becdelievre, Corine Gartioux, Valérie Jobic, et al.. Clinical and Molecular Spectrum Associated with COL6A3 c.7447A>G p.(Lys2483Glu) Variant: Elucidating its Role in Collagen VI-related Myopathies. *Journal of Neuromuscular Diseases*, 2021, 8 (4), pp.633-645. 10.3233/JND-200577 . hal-03244986

HAL Id: hal-03244986

<https://hal.u-pec.fr/hal-03244986v1>

Submitted on 17 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Clinical and Molecular Spectrum Associated with *COL6A3* c.7447A>G p.(Lys2483Glu) Variant: Elucidating its Role in Collagen VI-related Myopathies

Rocío N. Villar-Quiles^{a,b}, Sandra Donkevoort^c, Alix de Becdelièvre^{d,e}, Corine Gartioux^b,
Valérie Jobic^c, Reghan Foley^c, Riley McCarty^c, Ying Hu^c, Rita Menassa^f, Laurence Michelf^f,
Gaelle Gousse^g, Arnaud Lacour^h, Philippe Petiotⁱ, Nathalie Streichenberger^j, Ariane Choumert^k,
Léa Declerck^k, J.A Urtizbera^l, Guilhem Sole^m, Alain Furbyⁿ, Matthieu Cerino^o, Martin Krahn^o,
Emmanuelle Campana- Salort^p, Ana Ferreira^{a,q}, Bruno Eymard^a, Carsten G. Bönnemann^b,
Pascale Richard^c, Valérie Allamand^{b,r}, Corinne Métay^{e,b,l} and Tanya Stojkovic^{a,b,l,*}

^aAP-HP, Reference Center for Neuromuscular Disorders, Pitié-Salpêtrière Hospital, Paris, France

^bSorbonne Université - Inserm UMRS974, Centre de Recherche en Myologie, GH Pitié-Salpêtrière, Paris, France

^cNeuromuscular and Neurogenetic Disorders of Childhood Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA

^dAP-HP, Département de Génétique, Hôpital Henri Mondor, Créteil, France

^eAP-HP, Centre de Génétique Moléculaire et Chromosomique, UF Cardiogénétique et Myogénétique Moléculaire et Cellulaire, GH Pitié-Salpêtrière, Paris, France

^fHospices Civils de Lyon, LBMMS, Service Biochimie Biologie Moléculaire Grand Est, Groupement Hospitalier Est, CBPE, Bron, France

^gService de Neuropédiatrie, CHU Saint Étienne, France

^hService de Neurologie, CHU Saint Étienne, France

ⁱNeurologie et Explorations Fonctionnelles Neurologiques, Centre de Référence Maladies Neuromusculaires de la Région Rhône-Alpes Hôpital de la Croix-Rousse, Lyon, France

^jHospices Civils de Lyon, Centre de Pathologie et Neuropathologie Est, Groupement Hospitalier Est, Bron, France

^kCentre des Maladies Rares Neurologiques, CHU Sud Réunion, France

^lHôpital Marin, Centre de Compétence Neuromusculaire, Hendaye, France

^mCentre de Référence des Maladies Neuromusculaires AOC, Hôpital Pellegrin, CHU Bordeaux, France

ⁿCentre de Référence des Maladies Neuromusculaires Rares Rhône-Alpes, Hôpital Nord, CHU de Saint-Etienne, France

^oAP-HM, Département de Génétique Médicale, Hôpital Timone Enfants, Assistance Publique Hôpitaux de Marseille, France

^lLast co-authors.

*Correspondence to: Dr Tanya Stojkovic, Centre de Référence des Maladies Neuromusculaires Nord / Est / Ile-de-France, Groupe

Hospitalier Pitié-Salpêtrière, 47–83 Boulevard de l'Hôpital, 75651 Paris Cedex 13, France. E-mail: tanya.stojkovic@aphp.fr.

^P*AAP-HM, Centre de Référence des Maladies Neuromusculaires, Hôpital Timone, Marseille, France*

^Q*Basic and Translational Myology Lab, UMR8251, University Paris Diderot/CNRS, Paris, France*

^r*Unit of Muscle Biology, Department of Experimental Medical Science, Lund University, Lund, Sweden*

Abstract.

Background: Dominant and recessive autosomal pathogenic variants in the three major genes (*COL6A1-A2-A3*) encoding the extracellular matrix protein collagen VI underlie a group of myopathies ranging from early-onset severe conditions (Ullrich congenital muscular dystrophy) to milder forms maintaining independent ambulation (Bethlem myopathy). Diagnosis is based on the combination of clinical presentation, muscle MRI, muscle biopsy, analysis of collagen VI secretion, and *COL6A1-A2-A3* genetic analysis, the interpretation of which can be challenging.

Objective: To refine the phenotypical spectrum associated with the frequent *COL6A3* missense variant c.7447A>G (p.Lys2483Glu).

Methods: We report the clinical and molecular findings in 16 patients: 12 carrying this variant in compound heterozygosity with another *COL6A3* variant, and four homozygous.

Results: Patients carrying this variant in compound heterozygosity with a truncating *COL6A3*-variant exhibit a phenotype consistent with COL6-related myopathies (COL6-RM), with joint contractures, proximal weakness and skin abnormalities. All remain ambulant in adulthood and only three have mild respiratory involvement. Most show typical muscle MRI findings. In five patients, reduced collagen VI secretion was observed in skin fibroblasts cultures. All tested parents were healthy heterozygous carriers. Conversely, two out of four homozygous patients did not present with the classical COL6-RM clinical and imaging findings. Collagen VI immunolabelling on cultured fibroblasts revealed rather normal secretion in one and reduced secretion in another. Muscle biopsy from one homozygous patient showed myofibrillar disorganization and rimmed vacuoles.

Conclusions: In light of our results, we postulate that the *COL6A3* variant c.7447A>G may act as a modulator of the clinical phenotype. Thus, in patients with a typical COL6-RM phenotype, a second variant must be thoroughly searched for, while for patients with atypical phenotypes further investigations should be conducted to exclude alternative causes.

Keywords: Collagen VI-related myopathies, *COL6A3*, collagen type VI, neuromuscular disorders, limb-girdle muscular dystrophy (LGMD), congenital muscular dystrophy (CMD), muscular MRI, NGS

INTRODUCTION

Collagen VI is an extracellular matrix protein present in most tissues, notably in muscle, skin, tendon and blood vessels. Dominant or recessive autosomal mutations in each of the three “major” genes encoding the collagen VI α -chains (*COL6A1*, *COL6A2*, *COL6A3*) underlie collagen VI-related myopathies (COL6-RM), a heterogeneous group of disorders marked by a combined muscle and connective tissue involvement including joint laxity and contractures, as well as characteristic cutaneous abnormalities (*i.e.* keloid scars, keratosis pilaris and soft/velvety skin) in addition to muscle weakness.

COL6-RM clinical spectrum ranges from early-onset severe conditions (Ullrich congenital muscular dystrophy, UCMD) through phenotypes of intermediate severity to milder forms (Bethlem myopathy, BM) [1, 2]. Classically, UCMD patients present with congenital weakness and hypotonia, delayed motor milestones, associated with proximal joint contractures and concomitant marked distal hyperlaxity. Rigid spine, scoliosis, hip dislocation/dysplasia and prominent calcaneus are common features. Progressive

weakness leads to early loss of ambulation in most patients, and restrictive respiratory involvement occurs in most severely affected patients during the first two decades of life [3]. Conversely, BM phenotype is marked by milder proximal weakness associated with contractures typically affecting Achilles tendons, elbows, pectoralis, long finger flexors and interphalangeal joints. Although a slowly-progressive condition, two-thirds of patients over the age of 60 years may need assistance with ambulation [4, 5].

Nonetheless, there is a wide clinical variability, and intermediate phenotypes are now well recognized. Interestingly, patients can present with predominantly proximal weakness and very few or absent contractures or distal hyperlaxity, more akin to a limb-girdle muscular dystrophy (LGMD) [6, 7]. Furthermore, recessive mutations in *COL6A2* have also been associated with severe and widespread contractures known as myosclerosis [8].

Diagnosis is based on the combination of clinical presentation supported by muscle MRI, muscle biopsy findings, immunohistochemical examination of collagen VI secretion and analysis of the *COL6A1-3* genes sequences. Characteristic muscle

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

MRI findings of COL6-RM include a peculiar fatty replacement starting around the fascia surrounding the muscle (“outside-in” picture) with the presence of the so-called “central-cloud or shadow” typically affecting the *rectus femoris* and *vastus lateralis* [9].

Although UCMD was initially described as an autosomal recessive condition, dominant, mostly *de novo* mutations, have also been identified. Conversely, rare autosomal recessive mutations have been reported in BM patients, although it is mostly inherited as a dominant condition [4, 10]. Interpretation of genetic variants can be challenging, since there are few mutational hotspots, and the *COL6A1-3* genes are highly polymorphic. Furthermore, the clinical consequence of the many *COL6A1-3* variants that affect residues of unknown impact on the heterotrimeric assembly of COLVI (in particular variants outside of the triple helical domain) may be difficult to interpret and to validate experimentally.

We report a total of 16 patients carrying the *COL6A3* missense variant c.7447A>G either as compound heterozygotes with another *COL6A3* variant (12 patients from 10 families) or in a homozygous state (4 unrelated patients). Upon extensive analysis of the clinical phenotype and ancillary tests including muscle imaging pattern, muscle biopsy histology and COLVI secretion in dermal fibroblasts, we discuss the potential pathogenicity of this variant.

MATERIAL AND METHODS

Patients

Through an international collaboration, we identified 16 patients from 14 families (Table 1) carrying the c.7447A>G variant in the *COL6A3* gene in heterozygosity with another *COL6A3* mutation ($n = 12$) or in homozygosity ($n = 4$). Clinical data and ancillary tests (including serum CK levels, EMG, muscle MRI and muscle biopsy) were retrospectively retrieved and analyzed. All patients were examined by at least one of the authors in specialized neuromuscular departments. Diagnostic skeletal muscle biopsies were obtained, processed for standard histological and immunochemical studies and fixed for electron microscopy as previously described [11].

Informed consent was obtained from all patients in agreement with local ethical committees and with the 1964 Helsinki declaration and its later amendments (NIH, National Institute of Neurological Disorders and Stroke (NINDS), Institutional Review Board (Protocol 12N0095))

Genetic analysis, bioinformatics analysis and variants interpretation

Details on genetic testing and bioinformatics analysis can be found in Supplemental data. Pathogenicity of variants was determined according to current ACMG guidelines [12]. Variants were filtered out according to their allele frequency ($\leq 1\%$) as reported in the GnomAD database (<http://gnomad.broadinstitute.org/>). We then evaluated each variant considering a review of the literature, the location of the variant in the gene and the resulting corresponding protein, the *in silico* prediction tools (Polyphen2, SIFT, GVDG and CADD for missense variants and SpliceSiteFinder-like, MaxEntScan, NNSPLICE, GeneSplicer and Human Splicing Finder for splicing variants) and functional studies when available. SuSpect method (<http://www.sbg.bio.ic.ac.uk/suspect/>) was also used for prediction. All variants considered as pathogenic and likely pathogenic have been confirmed by a second independent method (Sanger sequencing).

Collagen VI immunolabelling

Dermal fibroblasts from eight index patients and one control individual were cultured to confluency in DMEM (Gibco) supplemented with 10% FBS (Biosera), penicillin/streptomycin (5700U Pen/5700 µg Strep; Gibco), and 50 µg/ml L-Ascorbic acid-2-phosphate (Sigma). Fixed cells were immunostained using either the polyclonal antibody Ab6588 (Abcam) as initially described in [13], or an $\alpha 3(\text{VI})$ specific antibody (HPA010080; Sigma-Aldrich). For the latter, cells were fixed with cold methanol prior to immunostaining. Confocal imaging was performed on a Nikon Ti2 microscope equipped with a motorized stage and a Yokogawa CSU-W1 spinning disk head coupled with a Prime 95 sCMOS camera (Photometrics). Z-stacks were obtained using a 0.15 µm steps, with a 40x/1.30 NA oil-immersion objective. Images were acquired with the same exposure setting, using the Metamorph software (Molecular Devices), and subsequently analyzed using Fiji [14]. Stacks were merged and Z projections (Sum slices) were obtained.

RESULTS

Genetics

Next Generation Sequencing (NGS) allowed the identification of the c.7447A>G p.(Lys2483Glu)

Table 1

Summarized clinical findings. DMM: delayed motor milestones; F : female; FH : follicular hyperkeratosis; FVC: forced vital capacity;LL : lower limbs; M : male; m : months; N/A : non-applicable; UL : upper limbs; y : years

Patient/ Gender	Onset	First symptoms	Age at last visit (y)	Weakness	Loss of ambulation	Contractures	Hyperlaxity Spine	Rigid involvement	Respiratory features	Skin
1-II.1/ M	Childhood	Poor sports performance	39	Axial+Proximal (UL 3/4, LL 3-4/5)	No	Wrists, elbows, knees, ankles	No	No	Yes(FVC 72%)	No
1-II.2/ F	Childhood	Difficulty in running, abnormal gait	31	Axial+Proximal (3/5 UL, 4/5 LL)	No. Walks with cane since 15y	Finger flexors, elbows, ankles	Yes	Yes	No	FH, prominent scars
2-II.1/ M	Childhood	Poor sports performance	62	Axial+Proximal (3/5 UL, LL)	No. Walks with cane since adulthood	Elbows, finger flexors, wrists, ankles, knees	No	Yes	No	No
3-III.1/ M	Early- childhood	Waddling gait, falls	18	Axial+Proximal (3/5UL,LL)	No	Finger flexors, ankles	Yes	Yes	Yes(FVC 68%)	FH
4-II.2/ M	Childhood	Difficulty running, joint contractures	51	Axial+Proximal	No	Elbows, fingers, wrists, ankles	No	No	No	No
5-II.2/ M	Early- childhood	DMM, frequent falls, ankle contractures	21	Axial+Proximal (4/5 UL, LL)	No	Finger flexors, elbows, ankles	No	Yes	No	Velvety skin
6-II.1/ M	Early- childhood	Abnormal gait, toe walking	55	Axial+Proximal (3 +/5 UL, LL)	No. Walks with cane since adulthood	Fingers, elbows, knees, ankles	No	No	No	No
7-II.1/ M	Adulthood	Waddling gait, proximal weakness	64	Axial+Proximal (4/5 UL, LL)	No	Finger flexors, wrist flexors, triceps surae	No	No	No	No
8-II.2/ F	At birth	Congenital torticollis DMM, falls, waddling gait	3.5	Axial+Proximal (4/5 UL, LL)	No	Ankles	Yes	No	No	No
9-II.5/ M	Childhood	Proximal weakness	54	Axial+Proximal (4/5 UL, LL)	No	Finger flexors, elbows, ankles	No	No	Yes(FVC 60%)	FH
10-II.1/ F	At birth	Decreased fetal movements, DMM	7	Axial+Proximal (4/5UL, LL)+Distal (3-4/5)	No	Ankles	Yes	No	No	No
10-II.3/ M	Congenital	Decreased fetal movements, DMM	2	Axial+Proximal	No	Ankles	Yes	No	N/A	FH
11-II.5/ M	Childhood	Bilateral <i>equinovarus</i> steppage gait	49	Distal (3/5 LL)	No	No	No	No	No	No
12-II.2/ M	Early childhood	Foot deformities, toe-walking	13	Proximal	No	Ankles	No	No	No	No
13-II.1/ F	Childhood	<i>Pes cavus</i> , abnormal gait	60	Axial+Proximal	No	Ankles	No	No	No	No
14-II.1/ F	Early childhood	DMM, difficulty running	77	Proximal (3/5 UL, LL)+ Distal (4/5 LL)	No, walker- since 74y	Finger flexors, ankles	No	No	No	No

variant in 16 patients, 5 females and 11 males ranging from age 2 to 77 years (mean 37.6 ± 21.8) at last examination (Table 1).

The *COL6A3* c.7447A>G missense variant (rs139260335 in dbSNP) is located in exon 36. At the protein level, the substitution affects the C1/11th Von Willebrand factor A domain (VWA 11), located in

a non-helical domain, and leads to an amino acid change with no significant post-translational modifications.

This variant was found either in compound heterozygosity associated with another exonic deletion, truncating or splicing *COL6A3* variant ($n=12$) or in homozygous state ($n=4$) (Table 2). Segregation




Fig. 1. Family pedigrees. Black filled symbols: affected individual. Crossed symbol: deceased individual.

190 studies were performed in eight families and
 191 confirmed that the tested parents were healthy het-
 192 erozygous carriers (Fig. 1).

193 Four patients (11-II.5, 12-II.2, 13-II.1 and 14-
 194 II.1) were homozygous for the c.7447A>G *COL6A3*
 195 variant. Patient 13-II.1's daughter was a healthy het-
 196 erozygous carrier. No segregation studies could be
 197 performed for patients 11-II.5, 12-II.2 and 14-II.1.
 198 Patient 13-II.1 also harbored a heterozygous *TNXB*
 199 variant c.4535_4552del, p.(Asp1512_Val1517del),
 200 whose allelic frequency was found to be low in
 201 gnomAD (13 allele count/275252) and reported as
 202 probably pathogenic in one patient in the LOVD
 203 database (reference #0000528445) associated with a
 204 known pathogenic *TNXB* variant. No second *TNXB*
 205 variant was identified in this patient.

206 Furthermore, no other variant that could affect
 207 splice, were found in the *COL6A1-A2-A3* genes after
 208 sequencing of the coding regions from cultured skin
 209 fibroblasts for patients 12-II.2, 13-II.1 and 14-II.1
 210 harboring the c.7447A>G at the homozygous state
 211 (no skin biopsy was available for 11-II.5 patient) (data
 212 not shown). Whole exome sequencing (WES) was
 213 performed in three compound heterozygous patients
 214 (7-II.1, 8-II.2, 10-II.1), thereby excluding additional
 215 pathogenic variants in other disease-causing genes.

216 Regarding patient 11-II.5, NGS panel for analy-
 217 sis of genes related to LGMD, distal, myofibrillar
 218 panels including *NEB* and *ACTA1* genes (responsible
 219 for most forms of nemaline myopathy) disclosed no
 220 pathogenic variants.

221 *Clinical presentation of compound heterozygous* 222 *patients*

223 Eleven (92%) patients developed the first symp-
 224 toms at birth or in early childhood, one around pub-
 225 erty and one in adulthood. None of them had a
 226 family history of neuromuscular disease. The most
 227 commonly recognized presenting symptoms were
 228 delayed motor development, waddling gait, poor sports
 229 performance and joint contractures. Decreased
 230 fetal movements were noted in two patients. Gait
 231 acquisition was delayed in two patients (5-II.2 and
 232 10-II.3) who reached independent ambulation at 18
 233 and 21 months, respectively. Two additional patients
 234 (8-II.2 and 10-II.1) started walking at 14 months and
 235 were noted to have a waddling gait. Muscle weakness
 236 was present in all patients with predominant axial and
 237 proximal involvement in 11 (92%), MRC grade 3-
 238 4/5. Prominent distal weakness was observed in one
 239 patient, affecting finger flexors and extensors in the

240 upper limbs and ankle dorsiflexion and eversion in
 241 the lower limbs (MRC 3/5). Independent ambulation
 242 was maintained in all patients although three patients
 243 (25%) required assistance (Table 1).

244 Joint contractures were present in all patients,
 245 mainly affecting finger flexors, elbows and ankles.
 246 Rigid spine was noted in four patients (33%). None of
 247 them had scoliosis. Distal hyperlaxity was observed
 248 in four patients (33%). Five patients (42%) were
 249 found to have cutaneous abnormalities including
 250 hyperkeratosis, keloid scars and velvety skin.

251 Forced Vital Capacity was measured in 15 patients
 252 and showed mild restrictive respiratory insufficiency
 253 in three of them (1-II.1, 3-III.1 and 9-II.5) but they did
 254 not need ventilatory support (Table 1). One patient (4-
 255 II.2) required a pacemaker due to left bundle branch
 256 block at age 48 years.

257 Serum Creatine Kinase (CK) levels were mildly
 258 to moderately elevated in all patients (range 120-
 259 970 IU/L). Electromyography (EMG) was performed
 260 in 10 patients, which revealed predominately myo-
 261 pathic findings in seven patients. Patients 8-II.2 and
 262 10-II.1 were found to have normal nerve conduction
 263 studies (NCS). EMG was limited due to poor tol-
 264 erance in patient 8-II.2. For patient 10-II.1, EMG
 265 at age 5 years disclosed slightly increased duration
 266 and amplitude and mildly reduced recruitment of the
 267 left *vastus lateralis*. Lastly, for patient 9-II.5, NCS
 268 showed absent peroneal motor response recorded at
 269 the *extensor digitorum brevis* bilaterally and EMG
 270 showed small motor units suggestive of a myopathic
 271 pattern in the proximal upper limbs, but neurogenic
 272 changes of the distal and proximal legs at age 55
 273 years. Muscle MRI (Fig. 2A and Supplementary Fig-
 274 ure 1) was performed in eight patients and showed
 275 characteristic COL6-RM findings in seven, namely
 276 fatty replacement starting around the fascia surround-
 277 ing *vastus lateralis* ("outside in") and the so-called
 278 "central cloud" pattern in the *rectus femoris*. Two
 279 patients' MRI showed fatty replacement of pelvic
 280 girdle and proximal leg muscles without the charac-
 281 teristic previously mentioned findings. Eight patients
 282 underwent a muscle biopsy, revealing prominent dys-
 283 trophic features in seven (87.5%).

284 *Clinical findings and ancillary tests in* 285 *homozygous patients*

286 Recognition of first symptoms was in childhood in
 287 all four patients. Patient 11-II.5 presented with foot
 288 deformities and steppage gait first recognized at age
 289 10 years. Patient 12-II.2 displayed equinovarus feet

Table 2

Genetics and ancillary tests. COL6-RM: Collagen VI related myopathies; FSV: fiber size variation; IN: internalized nuclei; LL : lower limbs; N/A : non-applicable; NCS: nerve conduction studies; ns: not shown;UL: upper limbs; WB : Western Blot

Patient	COL6A3 variants	CK (IU/L)	Muscle biopsy	Muscle MRI	Collagen VI secretion	Compatible COL6-RM phenotype
1-II.1	c.7447A>G,p.Lys2483Glu+c.4789 C>T,p.Arg1597*	190	N/A	Typical COL6-RM features	N/A	Yes
1-II.2	c.7447A>G,p.Lys2483Glu+c.4789 C>T,p.Arg1597*	200	Dystrophic	Typical COL6-RM features	Reduced	Yes
2-II.1	c.7447A>G,p.Lys2483Glu +c.9045 dupC,p.Gly3016Argfs*6	185	Dystrophic	Typical COL6-RM features	Reduced (ns)	Yes
3-III.1	c.7447A>G,p.Lys2483Glu+c.9384_9386 delGTA, p.Trp3128_Tyr3129delinsCys	120	Dystrophic	Typical COL6-RM features	Reduced (ns)	Yes
4-II.2	c.7447A>G,p.Lys2483Glu +c.6538_6591del,p.Val2181_Gly2198del	700	Dystrophic	N/A	Reduced	Yes
5-II.2	c.7447A>G,p.Lys2483Glu +c.6733C>T,p.Gln2245*	500	Dystrophic	Fatty replacement of pelvic girdle and proximal LL muscles	N/A	Yes
6-II.1	c.7447A>G,p.Lys2483Glu+c.7024C>T,p.Arg2342*	219	Dystrophic	Typical COL6-RM features	Reduced	Yes
7-II.1	c.7447A>G,p.Lys2483Glu+c.9329-1G>T	600	Dystrophic	Typical COL6-RM features	N/A	Yes
8-II.2	c.7447A>G,p.Lys2483Glu+c.6181C>T, p.Arg2061*	470	N/A	N/A	N/A	Yes
9-II.5	c.7447A>G, p.Lys2483Glu+c.7024C>T, p.Asg2342*	500-900	Fibre atrophy without necrosis	Typical COL6-RM features	N/A	Yes
10-II.1	exon 12 deletion [arr[GRCh37] 2q37.3 (238274212_238274975)x1]+c.7447A>G, p.Lys2483Glu	250	N/A	N/A	N/A	Yes
10-II.3	exon 12 deletion [arr[GRCh37] 2q37.3 (238274212_238274975)x1]+c.7447A>G, p.Lys2483Glu	170	N/A	N/A	N/A	Yes
11-II.5	c.7447A>G,p.Lys2483Glu homozygous	700	Mild FSV, IN, myofibrillar disorganization, autophagic vacuoles and nemalin rods	Atrophy and fatty infiltration of distal LL muscles (<i>gastrocnemius</i>), <i>peroneus lateralis</i>	N/A	No
12-II.2	c.7447A>G,p.Lys2483Glu homozygous	1600-2000	Dystrophic. Diminished calpain 3 expression on WB	Mild fatty infiltration of proximal LL muscles (<i>glutei</i> , <i>iliopsoas</i> , <i>quadriceps</i>)	N/A	No
13-II.1	c.7447A>G,p.Lys2483Glu homozygous	200	Dystrophic	Typical COL6-RM features	Normal	Yes
14-II.1	c.7447A>G,p.Lys2483Glu homozygous	157	FSV, muscle fiber atrophy without necrosis or regeneration	Typical COL6-RM features	Reduced (ns)	Yes




Fig. 2. A. Muscle imaging. Axial T1-weighted images. MRI from patients 1-II.1 (a) 6-II.1 (b) and 13-II.1 (c) show typical COL6-RM radiological findings, including fatty replacement starting around the fascia surrounding the muscle and the so-called “central-cloud” affecting the rectus femoris. (d,e) MRI from patient 11-II.5 reveals atrophy and fatty infiltration of distal lower limb muscles mainly affecting medial gastrocnemius and peroneus. (f) MRI from patient 12-II.2 showed no major muscle atrophy and mild fatty replacement of anterior and posterior thigh muscles. B. Muscle biopsy from patient 11-II.5. Hematoxylin and eosin (a) transversal frozen section show mild fiber size variation, numerous internalized nuclei and few rimmed vacuoles (arrow), while NADH-TR (b) reveal diffuse areas lacking oxydative activity. Note slightly irregular or discontinuous COLVI immunostaining (c). (d) Myofibrillar disorganization, autophagic vacuoles (arrow-d) and nemalin rods (arrow-e) are observed by electron microscopy (d,e). (f) COLVI and perlecan co-immunostaining. Scale bars: (a-c,f) 10 μ m; (d) 15 μ m; (e) 1 μ m.

290 and toe-walking since early-childhood, patient 13-
 291 II.1 presented with abnormal gait since childhood and
 292 patient 14-II.1 presented since early childhood with
 293 delayed motor milestones (delayed gait acquisition)
 294 and difficulty running. The latter three patients were
 295 found to have proximal weakness, while patient 11-
 296 II.5 had predominant distal weakness and bilateral

297 *scapula alata*. Patient 14-II.1 had finger flexor and
 298 Achilles contractures. Joint contractures were absent
 299 in the remaining three patients, aside from mild ankle
 300 contractures in patients 12-II.2 and 13-II.1 and none
 301 of them had distal hyperlaxity, skin abnormalities
 302 or respiratory involvement. Cardiac examination was
 303 normal in patients 11-II.5, 13-II.1 and 14-II.1, while

304 echocardiogram from patient 12-II.2 revealed mild
 305 aortic insufficiency at age 13 years. All patients main-
 306 tained ambulation but patient 14-II.1, currently 77
 307 years-old, requires assistance (walker) since age 74.

308 CK levels were strikingly elevated in patients 11-
 309 II.5 and 12-II.2, ranging from 700 to 2000 IU/L, while
 310 patients 13-II.1 and 14-II.1 had normal CK levels.
 311 Muscle MRI from the first two patients revealed non-
 312 typical COL6-RM findings (Fig. 2A). Indeed, fatty
 313 infiltration of posterior compartment of distal lower
 314 limb muscles was observed in patient 11-II.5 and
 315 mild fatty infiltration of proximal limb muscles was
 316 noticed in patient 12-II.2. Muscle MRI from patients
 317 13-II.1 and 14-II.1 showed typical COL6-RM find-
 318 ings affecting *vastus lateralis* and *rectus femoris*.

319 Muscle biopsy from patient 11-II.5 showed inter-
 320 nalized nuclei, myofibrillar disorganization, autoph-
 321 agic vacuoles and nemaline rods (Fig. 2B), while
 322 dystrophic changes were observed for patients 12-
 323 II.2 and 13-II.1 and muscle biopsy from patient
 324 14-II.1 disclosed only mild atrophy without necrosis
 325 or regeneration and fiber size variation.

326 Collagen secretion in cultured skin fibroblasts

327 Collagen VI immunolabelling was performed us-
 328 ing two different antibodies, on fixed dermal fibro-
 329 blasts derived from five compound heterozygous
 330 patients (Table 2, Fig. 3 and Supplementary Figure 2),
 331 and revealed reduced collagen VI secretion compared
 332 to control cells, in which a dense network of deposited
 333 collagen VI was detected. Since patients 1-II.2, 2-II.1
 334 and 6-II.1 harbor a second mutation introducing a
 335 premature termination codon that should lead to tran-
 336 script degradation *via* the nonsense-mediated decay,
 337 the COLVI staining detected most likely reflects the
 338 protein synthesis sustained by the missense-bearing
 339 allele. Similarly, the second mutation carried by pa-
 340 tients 3-III.1 and 4-II.2 (delins and exon delet-
 341 ion, respectively) should impair the assembly of
 342 monomers and/or secreted microfibrils. These results
 343 suggest that the missense variant does not signifi-
 344 cantly prevent COLVI assembly and secretion. Inter-
 345 estingly, COLVI immunolabelling of fibroblasts from
 346 3 homozygous patients revealed a rather normal
 347 secretion pattern (patients 11-II.5 and 13-II.1) or
 348 reduced secretion (14-II.1) (Fig. 3 and Supplemen-
 349 tary Figure 2). Accordingly, COLVI immunostaining
 350 on muscle biopsy from patient 11-II.5 was also rather
 351 preserved, with only focal points of discontinuous
 signal (Fig. 2c and 2f).




Fig. 3. Confocal imaging of COLVI secretion in fixed dermal fibroblasts from a control individual (CT) and 6 patients. COLVI (red) was detected with two polyclonal antibodies: Ab6588 (Abcam) on non-permeabilized cells (left panel) or an $\alpha 3(VI)$ chain specific on methanol-fixed cells (right panel). Nuclei are identified by DAPI staining (blue). In three compound heterozygous patients (1-II.2, 3-III.1 and 4-II.2) reduced collagen VI secretion was observed. Immunolabelling of fibroblasts from homozygous patients 11-II.5 and 13-II.1 revealed a rather normal secretion, while it was clearly reduced in the culture from homozygous patient 14-II.1. Scale bars = 50 μ m.

DISCUSSION

COL6-RM diagnosis is based on the combination of clinical presentation (*i.e.* muscle weakness, prominent contractures, characteristic cutaneous abnormalities and variable respiratory involvement), muscular imaging, immunohistochemical examination of collagen VI on muscle biopsy and fibroblasts and analysis of the *COL6A1-3* genes.

Although the first mutations detected in UCMD were recessive null mutations, leading to an absence of collagen VI in muscle biopsy sections and in cultured dermal fibroblasts [15], *de novo* dominant mutations in the *COL6A1-3* genes are responsible for a large proportion of UCMD cases [1, 16–18]. In BM, most patients harbor heterozygous dominantly acting mutations, typically affecting glycine residues of the Gly-X-Y motif at the N-terminal end of the triple helical domain that exert a dominant-negative effect on the tetramer structure [6, 17, 19–21]. Nonetheless, recessive mutations in the *COL6A1-3* genes have also been detected in patients with typical BM [22, 23], most of them carrying a null mutation on one allele in heterozygosity with a missense mutation on the other.

Interpretation of genetic variants can be challenging and many variants in the *COL6A1-3* genes have not yet been fully characterized at the functional level. Such is the case of missense variants affecting regions outside of the triple helix. Furthermore, there is a number of patients with a compatible clinical and muscle imaging phenotype who have no detectable mutations in the three *COL6* genes coding sequence [24, 25]. In that sense, 5′/3′ UTR regulatory elements or deep intronic splice mutations can go undetected by standard sequencing approaches, and muscle or fibroblasts RNA sequencing can be useful to detect these pathogenic variants [26].

Along these lines, the interpretation of the *COL6A3* c.7447A>G variant is complex. Its allelic frequency is 171/277005 and one homozygous individual is reported in GnomAD database. It is most prevalent in the non-Finnish European (0.001) and Latino (0.0007) populations. Its high allelic frequency raises questions about its pathogenicity, which remains unclear so far. Its predicted effect on SuSPect method (<http://www.sbg.bio.ic.ac.uk/suspect/>) [27] points to a low pathogenicity score (8 out of 100). The CADD score for this variant is 22.5 and it is classified as likely pathogenic or variant of unknown-signification in LOVD, Clinvar and likely pathogenic in the HGMDPro-database.

We report twelve patients carrying the c.7447A>G *COL6A3* variant in compound heterozygosity with a second *COL6A3* mutation. Most of these patients presented since childhood with proximal weakness associated with joint contractures, variable presence of rigid spine and skin abnormalities. All remained ambulatory at a mean age of 34.7 ± 21.5 and mild respiratory involvement was detected in three of them. This phenotype would be consistent with Bethlem myopathy (BM) [4, 7]. Muscle MRI showed typical COL6-RM findings in seven out of eight patients and collagen VI deposition in the extracellular matrix was reduced in dermal fibroblasts from five patients. WES analysis in three compound heterozygous patients excluded additional pathogenic variants in other neuromuscular disease-causing genes.

Conversely, homozygous patients exhibit strikingly different clinical features. Two out of four did not have a typical COL6-RM phenotype and presented with foot deformities, wing scapula and distal weakness or abnormal gait with toe-walking without major contractures, distal hyperlaxity, skin abnormalities or respiratory involvement. Moreover, CK levels of these two patients were strikingly high for COL6-RM, muscle MRI disclosed non-specific findings including only distal involvement in one, and muscle biopsy from the one patient showed myofibrillar disorganization, autophagic vacuoles and nemalin rods. Splicing defects were excluded after sequencing of the *COL6A1-A2-A3* coding regions in three patients harboring c.7447G>A at the homozygous state.

The *COL6A3* c.7447A>G variant has been previously reported in compound heterozygous patients [5, 28–31] with a clinical spectrum ranging from a mild phenotype when associated with a missense variant [30], to an intermediate phenotype with childhood onset and respiratory involvement [5] and a severe Ullrich-like phenotype [29], when associated with a second truncating *COL6A3* variant. To our knowledge, six homozygous patients have been reported so far [5, 29, 32]. Panadés-de Oliveira *et al.* [5] reported two homozygous siblings. The index case had proximal weakness, elevated CK levels (1000 IU/L) and dystrophic findings and rimmed vacuoles on muscle biopsy. The patient's sibling had asymptomatic significantly increased CK levels (4000 IU/L) detected at mid-age. Both had typical COL6-RM findings on muscle MRI findings. This study reported one additional homozygous patient who also carried a *COL6A1* c.2435-2A>G pathogenic splicing variant [5]. Interestingly, rimmed vacuoles were also found on the muscle biopsy from our patient 11-II.5.

454 The patient reported by Hunter et al. [29], pre-
455 sented with club feet, abnormal gait and lipoatrophy,
456 developed scoliosis and hamstrings, ankles and feet
457 contractures with overlapping toes. EMG studies
458 revealed a chronic motor neuropathy and muscle
459 biopsy disclosed abnormal myofibrillar architecture
460 with fiber type grouping. Recently, Stavusis et al.
461 [32] reported two siblings with mild proximal weak-
462 ness and joint contractures carrying three *COL6A3*
463 variants: the c.7447A>G variant in homozygosity
464 combined with the heterozygous frameshift variant
465 c.8074delT, p.(Tyr2692MetfsTer15). Strikingly, they
466 also reported a third patient, carrying the *COL6A3*
467 c.7447A>G variant in homozygosity, with proxi-
468 mal and distal weakness, diminished ankle reflexes
469 and axonal polyneuropathy, but no further biologi-
470 cal or genetic analysis regarding this polyneuropathy
471 is reported. Interestingly, collagen VI is expressed
472 in peripheral nerve [33–35]. Nonetheless, *COL6A1-*
473 *A2-A3* mutations have not been found associated
474 with hereditary neuropathies. On another note, mus-
475 cle MRI of this patient showed diffuse and severe
476 fatty infiltration without typical COL6-RM findings
477 and CK levels were also elevated from two to four
478 times the normal values. Unfortunately, no collagen
479 VI immunolabeling on muscle biopsy or analysis of
480 collagen VI production in dermal fibroblast cultures
481 are reported for any of these homozygous patients.

482 Stavusis et al. [32] have also reviewed all the rep-
483 orted patients to date, and conclude that for com-
484 pound heterozygous patients, the phenotype severity
485 entirely depends on the second mutation. The authors
486 speculate that this variant could perturb the bind-
487 ing properties of the protein and thus the stability of
488 the heterotrimer and modulate the phenotype when
489 present along with other mutations, as is the case for
490 the patients with the c.8074delT variant, which leads
491 to a premature termination of the protein. Nonethe-
492 less, the occurrence of another deep intronic mutation
493 in the *COL6A1-3* genes cannot be excluded [32].
494 Strikingly, reported homozygous patients exhibit
495 different clinical phenotypes such as peripheral neu-
496 ropathy [29, 32], asymptomatic hyperCK [5] or
497 limb-girdle myopathy with joint contractures [5, 32],
498 associated in some cases with a non-typical COL6-
499 RM histopathological presentation including rimmed
500 vacuoles [32] or abnormal myofibrillar architecture
501 with fiber type grouping [29].

502 At the functional level, the determinant role of
503 the $\alpha 3(\text{VI})$ chain in the monomer assembly is well
504 established [36]. However, the exact contribution of
505 the VWA C1 domain, harboring the mutated Lysine

506 residue at position 2483, remains poorly understood,
507 although it has been shown to be important for chain
508 recognition and assembly of triple helical molecules
509 [37]. Immunostaining of muscle biopsy and fibrob-
510 last cultures show that the missense variant does not
511 prevent COLVI deposition in the ECM, although fur-
512 ther analyses, such as Western blotting, should be
513 performed to obtain quantitative data. VWA domains
514 being protein-protein interaction modules, the mis-
515 sense variant may alter interaction with binding
516 partners as suggested by Stavusis et al. [32] but further
517 investigations are needed to support this hypothesis
518 experimentally.

519 In conclusion, this work expands the clinical and
520 molecular spectrum of COLVI-related myopathies
521 and is focused on the complex interpretation of the
522 frequent *COL6A3* c.7447A>G variant and the asso-
523 ciated clinical findings. Its high allele frequency and
524 the different phenotypes observed in the homozy-
525 gous patients reported so far, raise questions about
526 the pathogenicity of this variant in homozygosity. Our
527 results suggest that the *COL6A3* variant c.7447A>G
528 may act as a modulator of the clinical phenotype.
529 An in-depth analysis of clinical features and ancil-
530 lary tests is mandatory in order to interpret the genetic
531 analysis. For homozygous patients with a compatible
532 COL6-RM phenotype, including clinical, MRI find-
533 ings or altered collagen VI secretion, we recommend
534 to thoroughly search for an additional genetic variant
535 in any of the *COL6A1-3* genes, as in the cases reported
536 by Panades et al. [5] and Stavusis et al. [32], includ-
537 ing deep intronic variants leading to aberrant splicing
538 that may go undetected in NGS based panels [26].
539 Thus, analysis of mRNA transcripts could be use-
540 ful. In the event of a non-compatible phenotype such
541 as atypical clinical findings, strikingly elevated CK
542 levels, non-pathognomonic COL6-RM muscle MRI
543 findings, further genetic analysis would be advis-
544 able to exclude alternative causes such as inherited
545 neuropathies or other forms of myopathies. Further
546 analysis of homozygous carriers with detailed clini-
547 cal and ancillary tests data together with molecular
548 and genetic studies would help to elucidate whether
549 this variant is disease-causing or not. This would be of
550 major importance to allow precise diagnosis, accurate
551 management and genetic counseling for the affected
552 patients and families.

553 ACKNOWLEDGMENTS

554 We thank the patients and their families for par-
555 ticipating in our research study and Christopher

556 Mendoza, Christine Jones, and Gilberto (“Mike”)
 557 Averion for their help in the clinical setting. The
 558 (monoclonal antibody HFN7.1) developed by The
 559 University of Texas Health Science Center at San
 560 Antonio was obtained from the Developmental Stud-
 561 ies Hybridoma Bank, created by the NICHD of the
 562 NIH and maintained at The University of Iowa,
 563 Department of Biology, Iowa City, IA 52242.

564 FUNDING

565 The work in C.G. Bönnemann’s laboratory is sup-
 566 ported by intramural funds from the NIH National
 567 Institute of Neurological Disorders and Stroke. V.
 568 Allamand’s group is supported by intramural funds
 569 from the Association Institut de Myologie, and
 570 Inserm.

571 CONFLICT OF INTEREST

572 The authors have no conflict of interest to report.

573 REFERENCES

574 [1] Bushby KMD, Collins J, Hicks D. Collagen Type
 575 VI Myopathies. *Adv Exp Med Biol.* 2014;802:185-99.
 576 doi:10.1007/978-94-007-7893-1_12

577 [2] Bönnemann CG. The collagen VI-related myopathies.
 578 *Handb Clin Neurol.* 2011;101:81-96. doi:10.1016/B978-0-
 579 08-045031-5.00005-0

580 [3] Yonekawa T, Nishino I. Ullrich congenital muscular
 581 dystrophy: Clinicopathological features, natural history
 582 and pathomechanism(s). *J Neurol Neurosurg Psychiatry.*
 583 2015;86:280-7. doi:10.1136/jnnp-2013-307052

584 [4] Deconinck N, Richard P, Allamand V, Behin A, Lafôret P,
 585 Ferreira A, et al. Bethlem myopathy: Long-term follow-up
 586 identifies COL6 mutations predicting severe clinical evo-
 587 lution. *J Neurol Neurosurg Psychiatry.* 2015;86:1337-46.
 588 doi:10.1136/jnnp-2013-307245

589 [5] Panadés-de Oliveira L, Rodríguez-López C, Cantero
 590 Montenegro D, Marcos Toledano M del M, Fernández-
 591 Marmiesse A, Esteban Pérez J, et al. Bethlem myopathy:
 592 A series of 16 patients and description of seven new asso-
 593 ciated mutations. *J Neurol.* 2019;266:934-41. doi:10.1007/
 594 s00415-019-09217-z

595 [6] Scacheri PC, Gillanders EM, Subramony SH, Veda-
 596 narayanan V, Crowe CA, Thakore N, et al. Novel
 597 mutations in collagen VI genes: Expansion of the Bethlem
 598 myopathy phenotype. *Neurology.* 2002;58:593-602. doi:10.
 599 1212/WNL.58.4.593

600 [7] Bönnemann CG. The collagen VI-related myopathies: Mus-
 601 cle meets its matrix. *Nat Rev Neurol.* 2011;7:379-90.
 602 doi:10.1038/nrneurol.2011.81

603 [8] Merlini L, Martoni E, Grumati P, Sabatelli P, Squarzone S,
 604 Urciuolo A, et al. Autosomal recessive myosclerosis myopa-
 605 thy is a collagen VI disorder. *Neurology.* 2008;71:1245-53.
 606 doi:10.1212/01.wnl.0000327611.01687.5e

[9] Mercuri E, Lampe A, Allsop J, Knight R, Pane M, Kinali M, 607
 et al. Muscle MRI in Ullrich congenital muscular dystrophy 608
 and Bethlem myopathy. *Neuromuscul Disord.* 2005;15:303- 609
 10. doi:10.1016/j.nmd.2005.01.004 610

[10] Kim SY, Kim WJ, Kim H, Choi SA, Lee JS, Cho A, 611
 et al. Collagen VI-related myopathy: Expanding the clin- 612
 ical and genetic spectrum. *Muscle Nerve.* 2018. doi:10. 613
 1002/mus.26093 614

[11] Dubowitz V, Sewry CA. *Muscle Biopsy: A Practical 615
 Approach.* Philadelphia: Saunders Elsevier; 2007. 616

[12] Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster 617
 J, et al. Standards and guidelines for the interpretation of 618
 sequence variants: A joint consensus recommendation of 619
 the American College of Medical Genetics and Genomics 620
 and the Association for Molecular Pathology. *Genet Med.* 621
 2015;17:405-24. doi:10.1038/gim.2015.30 622

[13] Hicks D, Lampe AK, Barresi R, Charlton R, Fiorillo 623
 C, Bonnemann CG, et al. A refined diagnostic algo- 624
 rithm for Bethlem myopathy. *Neurology.* 2008;70:1192-9. 625
 doi:10.1212/01.wnl.0000307749.66438.6d 626

[14] Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Long- 627
 gair M, Pietzsch T, et al. Fiji: An open-source platform 628
 for biological-image analysis. *Nat Methods.* 2012;9:676-82. 629
 doi:10.1038/nmeth.2019 630

[15] Vanegas OC, Bertini E, Zhang RZ, Petrini S, Minosse 631
 C, Sabatelli P, et al. Ullrich scleroatonic muscular dys- 632
 trophy is caused by recessive mutations in collagen 633
 type VI. *Proc Natl Acad Sci U S A.* 2001;98:7516-21. 634
 doi:10.1073/pnas.121027598 635

[16] Dominant collagen VI mutations are a common cause of Ull- 636
 rich congenital muscular dystrophy. - PubMed - NCBI n.d. 637
[https://www.ncbi.nlm.nih.gov/pubmed/?term=baker+2005](https://www.ncbi.nlm.nih.gov/pubmed/?term=baker+2005+collagen+Vi) 638
[+collagen+Vi](https://www.ncbi.nlm.nih.gov/pubmed/?term=baker+2005+collagen+Vi) (accessed April 22, 2020). 639

[17] Lampe AK, Bushby KMD. Collagen VI related mus- 640
 cle disorders. *J Med Genet.* 2005;42:673-85. doi:10.1136/ 641
 jmg.2002.002311 642

[18] Lampe AK, Zou Y, Sudano D, O’Brien KK, Hicks D, Laval 643
 SH, et al. Exon skipping mutations in collagen VI are 644
 common and are predictive for severity and inheri- 645
 tance. *Hum Mutat.* 2008;29:809-22. doi:10.1002/humu. 646
 20704 647

[19] Jöbssis GJ, Keizers H, Vreijling JP, Visser M de, Speer 648
 MC, Wolterman RA, et al. Type VI collagen mutations 649
 in Bethlem myopathy, an autosomal dominant myopa- 650
 thy with contractures. *Nat Genet.* 1996;14:113-5. doi:10. 651
 1038/ng0996-113 652

[20] Pepe G, Giusti B, Bertini E, Brunelli T, Saitta B, Comeglio 653
 P, et al. A heterozygous splice site mutation in COL6A1 654
 leading to an in-frame deletion of the $\alpha 1(VI)$ collagen 655
 chain in an Italian family affected by Bethlem myopa- 656
 thy. *Biochem Biophys Res Commun.* 1999;258:802-7. 657
 doi:10.1006/bbrc.1999.0680 658

[21] Lucieli S, Giusti B, Mercuri E, Vanegas OC, Lucarini 659
 L, Pietroni V, et al. Detection of common and private 660
 mutations in the COL6A1 gene of patients with Beth- 661
 lem myopathy. *Neurology.* 2005;64:1931-7. doi:10.1212/ 662
 01.WNL.0000163990.00057.66 663

[22] Foley AR, Hu Y, Zou Y, Columbus A, Shoffner J, Dunn 664
 DM, et al. Autosomal recessive inheritance of classic 665
 Bethlem myopathy. *Neuromuscul Disord.* 2009;19:813-7. 666
 doi:10.1016/j.nmd.2009.09.010 667

[23] Gualandi F, Urciuolo A, Martoni E, Sabatelli P, Squarzone 668
 S, Bovolenta M, et al. Autosomal recessive Beth- 669
 lem myopathy. *Neurology.* 2009;73:1883-91. doi:10.1212/ 670
 WNL.0b013e3181c3fd2a 671

- 672 [24] Tétrault M, Duquette A, Thiffault I, Bherer C, Jarry J, 707
673 Loisel L, et al. A new form of congenital muscular dys- 708
674 trophy with joint hyperlaxity maps to 3p23-21. *Brain*. 709
675 2006;129:2077-84. doi:10.1093/brain/awl146 710
- 676 [25] Petrini S, D'Amico A, Sale P, Lucarini L, Sabatelli P, 711
677 Tessa A, et al. Ullrich myopathy phenotype with secondary 712
678 ColVI defect identified by confocal imaging and electron 713
679 microscopy analysis. *Neuromuscul Disord*. 2007;17:587- 714
680 96. doi:10.1016/j.nmd.2007.04.010 715
- 681 [26] Bolduc V, Reghan Foley A, Solomon-Degefa H, Sarathy 716
682 A, Donkervoort S, Hu Y, et al. A recurrent COL6A1 717
683 pseudoexon insertion causes muscular dystrophy and is 718
684 effectively targeted by splice-correction therapies. *JCI* 719
685 *Insight*. 2019;4. doi:10.1172/jci.insight.124403 720
- 686 [27] Yates CM, Filippis I, Kelley LA, Sternberg MJE. SuS- 721
687 Pect: Enhanced prediction of single amino acid variant 722
688 (SAV) phenotype using network features. *J Mol Biol*. 723
689 2014;426:2692-701. doi:10.1016/j.jmb.2014.04.026 724
- 690 [28] Sframeli M, Sarkozy A, Bertoli M, Astrea G, Hudson J, 725
691 Scoto M, et al. Congenital muscular dystrophies in the 726
692 UK population: Clinical and molecular spectrum of a large 727
693 cohort diagnosed over a 12-year period. *Neuromuscul Dis-* 728
694 *ord*. 2017;27:793-803. doi:10.1016/j.nmd.2017.06.008 729
- 695 [29] Hunter JM, Ahearn ME, Balak CD, Liang WS, Kur- 730
696 doglu A, Corneveaux JJ, et al. Novel pathogenic variants 731
697 and genes for myopathies identified by whole exome 732
698 sequencing. *Mol Genet Genomic Med*. 2015;3:283-301. 733
699 doi:10.1002/mgg3.142 734
- 700 [30] Briñas L, Richard P, Quijano-Roy S, Gartioux C, Ledeuil 735
701 C, Lacène E, et al. Early onset collagen VI myopathies: 736
702 Genetic and clinical correlations. *Ann Neurol*. 2010;68:511- 737
703 20. doi:10.1002/ana.22087 738
- 704 [31] Fichna JP, Macías A, Piechota M, Korostynski M, 739
705 Potulska-Chromik A, Redowicz MJ, et al. Whole-exome 740
706 sequencing identifies novel pathogenic mutations and puta- 741
707 tive phenotype-influencing variants in Polish limb-girdle 742
708 muscular dystrophy patients. *Hum Genomics*. 2018;12. 743
709 doi:10.1186/s40246-018-0167-1 744
- [32] Stavakis J, Micule I, Wright NT, Straub V, Topf A, Panadés- 745
710 de Oliveira L, et al. Collagen VI-related limb-girdle syn- 746
711 drome caused by frequent mutation in COL6A3 gene with 747
712 conflicting reports of pathogenicity. *Neuromuscul* 748
713 *Disord*. 2020. doi:10.1016/j.nmd.2020.03.010 749
- [33] Chen P, Cescon M, Megighian A, Bonaldo P. Collagen 750
714 VI regulates peripheral nerve myelination and function. 751
715 *FASEB J*. 2014;28:1145-56. doi:10.1096/fj.13-239533 752
- [34] Chen P, Cescon M, Bonaldo P. The Role of Collagens in 753
716 Peripheral Nerve Myelination and Function. *Mol Neurobiol*. 754
717 2015;52:216-25. doi:10.1007/s12035-014-8862-y 755
- [35] Cescon M, Gregorio I, Eiber N, Borgia D, Fusto A, Sabatelli 756
718 P, et al. Collagen VI is required for the structural and 757
719 functional integrity of the neuromuscular junction. *Acta* 758
720 *Neuropathol*. 2018;136:483-99. doi:10.1007/s00401-018- 759
721 1860-9 760
- [36] Fitzgerald J, Mörgelin M, Selan C, Wiberg C, Keene DR, 761
722 Lamandé SR, et al. The N-terminal N5 subdomain of 762
723 the $\alpha 3(\text{VI})$ chain is important for collagen VI microfibril 763
724 formation. *J Biol Chem*. 2001;276:187-93. doi:10. 764
725 1074/jbc.M008173200 765
- [37] Lamandé SR, Mörgelin M, Adams NE, Selan C, Allen JM. 766
726 The C5 domain of the collagen VI $\alpha 3(\text{VI})$ chain is criti- 767
727 cal for extracellular microfibril formation and is present 768
728 in the extracellular matrix of cultured cells. *J Biol Chem*. 769
729 2006;281:16607-14. doi:10.1074/jbc.M510192200 770
- [38] Krahn M, Biancalana V, Cerino M, Perrin A, Michel- 771
730 Calemard L, Nectoux J, et al. A National French consensus 772
731 on gene lists for the diagnosis of myopathies using next- 773
732 generation sequencing. *Eur J Hum Genet*. 2019;27:349-52. 774
733 doi:10.1038/s41431-018-0305-1 775