



HAL
open science

Multicentre validation of a EUCAST method for the antifungal susceptibility testing of microconidia-forming dermatophytes

Maiken Cavling Arendrup, Karin Meinike Jørgensen, Jesus Guinea, Katrien Lagrou, Erja Chryssanthou, Marie-Pierre Hayette, Francesco Barchiesi, Cornelia Lass-Flörl, Petr Hamal, Eric Dannaoui, et al.

► To cite this version:

Maiken Cavling Arendrup, Karin Meinike Jørgensen, Jesus Guinea, Katrien Lagrou, Erja Chryssanthou, et al.. Multicentre validation of a EUCAST method for the antifungal susceptibility testing of microconidia-forming dermatophytes. *Journal of Antimicrobial Chemotherapy*, 2020, 75 (7), pp.1807-1819. 10.1093/JAC/DKAA111 . hal-03950328

HAL Id: hal-03950328

<https://hal.u-pec.fr/hal-03950328>

Submitted on 7 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ECCMID 2020 Abstract

Multicentre validation of a EUCAST method for the antifungal susceptibility testing of microconidia-forming Dermatophytes.

Maiken Cavling Arendrup^{1,2,3}, Karin Meinike Jørgensen¹, Jesus Guinea^{3,4}, Katrien Lagrou⁵, Erja Chryssanthou⁶, Marie-Pierre Hayette⁷, Francesco Barchiesi⁸, Cornelia Lass Flörl⁹, Petr Hamal¹⁰, Eric Dannaoui¹¹, Anuradha Chowdhary¹², Joseph Meletiadis¹³

Background: Terbinafine resistance is increasingly reported in *Trichophyton rubrum* and *Trichophyton interdigitale* rendering susceptibility testing important particularly in non-responding cases. We performed a multicentre evaluation of a recently proposed modified EUCAST method implementing medium supplemented with chloramphenicol and cycloheximide (CC) to avoid contamination.

Materials/methods: A blinded panel of wild-type and squalene epoxidase (SQLE) target gene mutant *T. rubrum* and *T. interdigitale* strains were distributed to 10 European laboratories. Susceptibility to terbinafine, itraconazole, voriconazole and amorolfine were performed according to the E.Def 9.3.1 method with and without addition of chloramphenicol and cycloheximide (final concentrations 50 mg/L and 300 mg/L, respectively). Plates were incubated at 25 °C (one laboratory used 30 °C) for 5-7 days until sufficient growth. MICs were determined visually (ignoring trailing growth for itraconazole) and spectrophotometrically with 90% and 50% endpoints yielding a total of 7,829 MICs. *A. flavus* ATCC 204304 and *A. flavus* CNM-CM1813 were included as controls.

Results: 100%/96% (voriconazole) and 84%/84% (itraconazole) MIC determinations fell within the QC ranges for the two QC strains, respectively, and 96%/92% terbinafine MICs fell in a 0.25-1 mg/L 3 two-fold-dilution range suggesting a high interlaboratory reproducibility. Across the six methods, the number of terbinafine MEs varied from 2 (2.6%) to 5 (6.6%) for *T. rubrum* and between 0 and 2 (2.0%) for *T. interdigitale* (lowest for the CC-method (2.6%-4.4%/ 0-1% for *T. rubrum*/*T. interdigitale*). The difference between the modes for the wt and mutant population were ≥ 7 two-fold-dilutions in all cases (Table). If excluding a I121M/V237I *T. rubrum* mutant, and two mixed *T. interdigitale* strains, the number of VMEs were CC visual: *T. rubrum*: 1/77 (1.3%), CC spec-90%: 3/68 (4.4%) and CC spec-50%: 1/76 (1.3%), and none for *T. interdigitale*. The activity of voriconazole, itraconazole and amorolfine were quite uniform against *T. rubrum* and *T. interdigitale*, but unacceptably wide MIC ranges were found for the visual and spec-90% inhibition methods for itraconazole (data not shown).

Conclusions: Although none of the laboratories perform dermatophyte testing at a regular basis an acceptable interlaboratory agreement and good separation between SQLE wt and mutants were found, suggesting a robust performance of the proposed method.