



HAL
open science

In Vitro Antifungal Combination of Flucytosine with Amphotericin B, Voriconazole, or Micafungin against *Candida auris* Shows No Antagonism

A. Bidaud, F. Botterel, A. Chowdhary, E. Dannaoui

► **To cite this version:**

A. Bidaud, F. Botterel, A. Chowdhary, E. Dannaoui. In Vitro Antifungal Combination of Flucytosine with Amphotericin B, Voriconazole, or Micafungin against *Candida auris* Shows No Antagonism. *Antimicrobial Agents and Chemotherapy*, 2019, 63 (12), 10.1128/AAC.01393-19 . hal-03997625

HAL Id: hal-03997625

<https://hal.u-pec.fr/hal-03997625v1>

Submitted on 4 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



In Vitro Antifungal Combination of Flucytosine with Amphotericin B, Voriconazole, or Micafungin against *Candida auris* Shows No Antagonism

A. L. Bidaud,^a  F. Botterel,^b  A. Chowdhary,^c  E. Dannaoui^{a,b}

^aParis-Descartes University, Faculty of Medicine, APHP, European Georges Pompidou Hospital, Parasitology-Mycology Unit, Microbiology Department, Paris, France

^bDynamic Research Group, Paris Est Créteil University (UPEC, EnvA), Créteil, France

^cDepartment of Medical Mycology, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi, India

ABSTRACT *Candida auris* is an emerging, multidrug-resistant pathogen responsible for invasive hospital-acquired infections. Flucytosine is an effective anti-*Candida* species drug, but which cannot be used as a monotherapy because of the risk of development of resistant mutants during treatment. It is, therefore, noteworthy to test possible combinations with flucytosine that may have a synergistic interaction. In this study, we determined the *in vitro* interaction between flucytosine and amphotericin B, micafungin, or voriconazole. These combinations have been tested against 15 *C. auris* isolates. The MIC ranges (geometric mean [Gmean]) of flucytosine, amphotericin B, micafungin, and voriconazole were 0.125 to 1 $\mu\text{g/ml}$ (0.42 $\mu\text{g/ml}$), 0.25 to 1 $\mu\text{g/ml}$ (0.66 $\mu\text{g/ml}$), 0.125 to 0.5 $\mu\text{g/ml}$ (0.3 $\mu\text{g/ml}$), and 0.03 to 4 $\mu\text{g/ml}$ (1.05 $\mu\text{g/ml}$), respectively. When tested in combination, indifferent interactions were mostly observed with fractional inhibitory concentration index values from 0.5 to 1, 0.31 to 1.01, and 0.5 to 1.06 for the combinations of flucytosine with amphotericin B, micafungin, and voriconazole, respectively. A synergy was observed for the strain CBS 10913 from Japan. No antagonism was observed for any combination. The combination of flucytosine with amphotericin B or micafungin may be relevant for the treatment of *C. auris* infections.

KEYWORDS *Candida auris*, antifungal, combination, amphotericin B, flucytosine, micafungin, voriconazole

Candida auris is a multidrug-resistant emerging fungal pathogen responsible for invasive infection (1–3). *C. auris* was first described from an external ear canal drainage specimen from a Japanese patient in 2009 (4). After 2009, *C. auris* was highlighted worldwide and has been the subject of many publications (5–9). But the cases of *C. auris* have been mainly reported from India, Central and North America, Spain, United Kingdom, Kuwait, South Africa, Israel, and Oman (10–16). Recently, an isolate of *C. auris* has been found in China (17), and the first case of *C. auris* infection has been described in France (18).

C. auris has a close phylogenetic relationship with *Candida haemulonii*, *Candida dubushaemulonii*, and *Clavispora lusitaniae* which can lead to erroneous identifications using commercial biochemical identification methods (19, 20). Matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) or molecular identification based on sequencing the D1-D2 region of the 28S ribosomal DNA allows the correct identification of *C. auris* (21).

The emergence of *C. auris* is alarming especially because this fungal pathogen has the potential to harbor or develop multidrug resistance. Only 4 drug classes are available for the systemic treatment of *Candida* infections, including the echinocandins (caspofungin, micafungin, and anidulafungin), azoles (fluconazole, itraconazole, vori-

Citation Bidaud AL, Botterel F, Chowdhary A, Dannaoui E. 2019. *In vitro* antifungal combination of flucytosine with amphotericin B, voriconazole, or micafungin against *Candida auris* shows no antagonism. *Antimicrob Agents Chemother* 63:e01393-19. <https://doi.org/10.1128/AAC.01393-19>.

Copyright © 2019 American Society for Microbiology. All Rights Reserved.

Address correspondence to E. Dannaoui, eric.dannaoui@aphp.fr.

Received 18 July 2019

Returned for modification 15 August 2019

Accepted 1 October 2019

Accepted manuscript posted online 7 October 2019

Published 21 November 2019

TABLE 1 *In vitro* interactions of amphotericin B with flucytosine against *Candida auris*^a

Isolate	MIC ($\mu\text{g/ml}$) of drug alone		MIC ($\mu\text{g/ml}$) of the drug in combination		Lowest FICI for the combination	Interaction
	AMB	5-FC	AMB	5-FC		
CBS 12372	0.25	0.5	0.06	0.25	0.75	N
CBS 12373	0.25	0.5	0.125	0.06	0.63	N
CBS 12766	1	0.5	0.5	0.06	0.63	N
CBS 12767	1	0.5	0.5	0.06	0.63	N
CBS 12768	1	0.5	0.5	0.06	0.63	N
CBS 12769	1	0.5	0.5	0.06	0.63	N
CBS 12770	0.5	0.5	0.25	0.25	1	N
CBS 12771	1	0.5	0.5	0.06	0.63	N
CBS 12772	1	0.5	0.5	0.06	0.63	N
CBS 12773	1	0.5	0.5	0.03	0.56	N
CBS 12774	1	0.5	0.5	0.06	0.63	N
CBS 12775	1	0.5	0.5	0.06	0.63	N
CBS 12776	1	0.5	0.5	0.06	0.63	N
CBS 12777	0.25	0.5	0.125	0.06	0.63	N
CBS 10913	0.25	0.25	0.06	0.06	0.50	S

^aAMB, amphotericin B; 5-FC, flucytosine; N, no interaction; S, synergy; FICI, fractional inhibitory concentration index.

conazole, posaconazole, and isavuconazole), polyenes (amphotericin B), and, finally, the pyrimidine analogue flucytosine (22). Resistance to each of these classes has been reported. Although uncommon, some *C. auris* strains exhibit elevated MICs for all three major classes of antifungal drugs, i.e., azoles, polyenes, and echinocandins (23, 24). There are currently recommendations for the treatment of *C. auris* infections with echinocandins as first-line therapy (25). Amphotericin B could be used for patients not responding to echinocandin therapy, depending on MIC results (2, 3). However, it has been shown that amphotericin B resistance is not uncommon (26).

Amphotericin B, flucytosine, echinocandins, and azoles have different cellular targets, and their combination might be of interest in the treatment of *C. auris* infections. The combination of flucytosine with amphotericin B has been shown *in vitro* to be synergistic against several fungal pathogens, such as *Cryptococcus neoformans* (27). Amphotericin B and flucytosine are used in several fungal infections, such as cryptococcosis, but also in *Candida* species infections (e.g., endocarditis, meningitis, and endophthalmitis) (28, 29). The hypothesis would be that amphotericin B, by forming pores of the fungal membrane, would facilitate the intracellular penetration of flucytosine. Furthermore, flucytosine also exhibits synergy with azoles against *C. neoformans* and *Candida albicans* both *in vitro* and *in vivo* (30–33).

The aim of the present study was to evaluate the *in vitro* interaction between flucytosine and either a polyene (amphotericin B), an echinocandin (micafungin), or an azole (voriconazole) against several *C. auris* isolates from different origins.

(This work was presented in part at ASM Microbe 2019, 20 to 24 June 2019, San Francisco, CA)

RESULTS

The results for the tested drugs alone and in combination against *C. auris* isolates are summarized in Tables 1, 2, and 3. All experiments were run in duplicate with similar results. MICs of the drugs alone were within ± 2 log₂ dilutions in 100% of the cases. Therefore, results from one replicate are shown. The MIC ranges of drugs alone against the strains were 0.125 to 1 $\mu\text{g/ml}$ for flucytosine, 0.25 to 1 $\mu\text{g/ml}$ for amphotericin B, 0.125 to 0.5 $\mu\text{g/ml}$ for micafungin, and 0.03 to 4 $\mu\text{g/ml}$ for voriconazole. With the checkerboard microdilution assay, when amphotericin was combined with flucytosine, the MIC ranges of amphotericin B and flucytosine decreased from 0.25 to 1 $\mu\text{g/ml}$ to 0.06 to 0.5 $\mu\text{g/ml}$ and from 0.125 to 1 $\mu\text{g/ml}$ to 0.03 to 0.25 $\mu\text{g/ml}$, respectively. These values correspond to a median (range) decrease of 1- (1- to 2-) fold and 3- (1- to 4-) fold for amphotericin B and

TABLE 2 *In vitro* interactions of micafungin with flucytosine against *Candida auris*^a

Isolate	MIC (µg/ml) of drug alone		MIC (µg/ml) of the drug in combination		Lowest FICI for the combination	Interaction
	MICA	5-FC	MICA	5-FC		
CBS 12372	0.25	0.5	0.125	0.008	0.52	N
CBS 12373	0.25	1	0.125	0.03	0.53	N
CBS 12766	0.5	0.5	0.25	0.008	0.52	N
CBS 12767	0.25	0.5	0.03	0.25	0.63	N
CBS 12768	0.5	0.5	0.25	0.008	0.52	N
CBS 12769	0.25	0.5	0.125	0.25	1	N
CBS 12770	0.25	0.25	0.125	0.125	1	N
CBS 12771	0.25	0.25	0.125	0.125	1	N
CBS 12772	0.5	0.25	0.25	0.015	0.56	N
CBS 12773	0.25	0.5	0.125	0.25	1	N
CBS 12774	0.25	0.5	0.25	0.008	1.02	N
CBS 12775	0.25	0.5	0.125	0.25	1	N
CBS 12776	0.5	0.5	0.25	0.008	0.52	N
CBS 12777	0.5	0.5	0.25	0.008	0.52	N
CBS 10913	0.125	0.125	0.03	0.008	0.31	S

^aMICA, micafungin; 5-FC, flucytosine; N, no interaction; S, synergy; FICI, fractional inhibitory concentration index.

flucytosine, respectively. The fractional inhibitory concentration index (FICI) ranged from 0.5 to 1 which was indicative of no-interaction except for one isolate (CBS 10913) for which a synergy was observed. When micafungin was combined with flucytosine, the MIC ranges of micafungin and flucytosine decreased from 0.125 to 0.5 µg/ml to 0.03 to 0.25 µg/ml and from 0.125 to 1 µg/ml to 0.008 to 0.25 µg/ml, respectively. This corresponds to a median (range) decrease of 1- (0- to 3-) fold and 4- (1- to 6-) fold for micafungin and flucytosine, respectively. Synergistic effects of micafungin with flucytosine were shown against the *C. auris* isolate number CBS 10913 from Japan (FICI, 0.31). When voriconazole was combined with flucytosine, the MIC ranges of voriconazole and flucytosine decreased from 0.03 to 4 µg/ml to 0.008 to 2 µg/ml and from 0.125 to 1 µg/ml to 0.008 to 0.5 µg/ml, respectively. These results correspond to a median (range) decrease of 7- (0- to 8-) fold and 1- (0- to 5-) fold for voriconazole and flucytosine, respectively. FICI ranged from 0.504 to 1.06, which was indicative of no interaction. No antagonist effects were observed for any combination.

An example of analysis (strain CBS 10913) with the response surface approach based on the Bliss model for the three combinations is shown in Fig. 1. Synergy was observed

TABLE 3 *In vitro* interactions of voriconazole with flucytosine against *Candida auris*^a

Isolate	MIC (µg/ml) of drug alone		MIC (µg/ml) of the drug in combination		Lowest FICI for the combination	Interaction
	VORI	5-FC	VORI	5-FC		
CBS 12372	1	0.25	0.5	0.125	1	N
CBS 12373	1	0.5	0.008	0.5	1	N
CBS 12766	1	0.5	0.03	0.25	0.53	N
CBS 12767	1	0.25	0.008	0.25	1	N
CBS 12768	1	0.5	0.008	0.25	0.51	N
CBS 12769	1	0.5	0.008	0.5	1	N
CBS 12770	4	0.25	2	0.008	0.53	N
CBS 12771	2	0.5	0.008	0.25	0.51	N
CBS 12772	1	0.25	0.008	0.25	1	N
CBS 12773	1	0.5	0.008	0.5	1	N
CBS 12774	2	0.5	0.008	0.5	1	N
CBS 12775	2	0.5	0.008	0.25	0.51	N
CBS 12776	1	0.5	0.008	0.25	0.51	N
CBS 12777	2	0.25	0.008	0.25	1	N
CBS 10913	0.03	0.125	0.03	0.008	1.06	N

^aVORI, Voriconazole; 5-FC, flucytosine; N, no interaction; S, synergy; FICI, fractional inhibitory concentration index.




FIG 1 Combination of flucytosine with amphotericin B (A), micafungin (B), and voriconazole (C) against *C. auris* CBS 10913 analyzed by response surface modeling. Combined results from two independent experiments were used for analysis.

for the combinations of flucytosine with either amphotericin B or micafungin, while the interaction between flucytosine and voriconazole was indifferent.

DISCUSSION

So far, evidence to support combination therapy in infections with *C. auris* is lacking, and clinicians make decisions on a case-by-case basis. Nevertheless, as *C. auris* is often a multidrug-resistant organism, combination therapy could be interesting, as suggested by the few *in vitro* studies performed (34–36). In a recent study, interactions between azoles and echinocandins showed synergy for the combination of micafungin with voriconazole against all the isolates tested (34). Moreover, the same study also showed indifferent interactions when micafungin was combined with fluconazole and indifferent interactions when caspofungin was combined with fluconazole or voriconazole.

In the present study, we tested the combination of flucytosine with a polyene, an echinocandin, or an azole. Flucytosine is an interesting anti-*Candida* species drug because this nucleic base analogue penetrates the fungal cells through a transporter (cytosine permease) and blocks DNA synthesis and protein synthesis. Flucytosine has excellent bioavailability *per os*. Because this molecule is water-soluble and not linked to plasma proteins, its distribution in most tissues is homogeneous as well as its diffusion in cerebrospinal fluid. Against *C. auris*, flucytosine is very active, as shown in the present study as well as in previous studies (21, 23, 37). Nevertheless, flucytosine is never used as a monotherapy because of the rapid selection of resistant mutants during treatment. Therefore, it seems advisable to test *in vitro* combinations with this molecule. The major interests of using a combination treatment with flucytosine are to benefit from the good flucytosine pharmacokinetics and to potentially have a synergistic interaction with the partner drug. A third motivation for combination therapy might be to reduce the risk of further resistance development in *C. auris* during therapy, although this is not trivial, as this organism has a high potential for resistance acquisition. It is also important to ensure that these drug combinations do not lead to antagonism.

In the present study, we used the checkerboard microdilution method. Analyzing results for this technique is not straightforward, particularly for the choice of the most suitable inhibition endpoint which could be different for analyzing the activity of the drugs alone or in combination. Moreover, interactions are only calculated for few concentrations (corresponding to the MIC). For this reason, we also used an alternative method of response surface modeling (based on the Bliss independency model) to visualize the interaction when synergy was found by the FICI method. The main advantages of this approach are that it is independent of any endpoint and it takes in account all the concentrations tested in the microplate. The same results were obtained by the two mathematical models used (Loewe and Bliss).

Overall, although the three combinations showed mainly indifferent interactions, we observed that flucytosine can be used in combination with other drugs without risk of antagonism. Of note, the only isolate against which synergy was observed was the type strain CBS 10913, characterized by its lower MICs than more recent isolates from India or other countries (23, 38). The synergistic interactions could be linked to this particular susceptibility profile. Nevertheless, despite the high susceptibility of CBS 10913 to voriconazole compared with the other strains, synergy was not observed for the voriconazole-flucytosine combination. One limitation of our study is the test of isolates only belonging to the East Asian or South Asian clades. No isolates representing other clades of *C. auris* were included. There are currently 4 or 5 phylogeographic clades (3, 37, 39). The clades are genetically and phenotypically distinct, and there could be alternative results among the other clades.

Results of antifungal combinations are important because therapeutic options for *C. auris* are limited. Although the echinocandins appear to be the best choice for initial treatment, the development of new antifungal drugs with activity against *C. auris* will be essential to control multidrug-resistant isolates. Before new drugs are available, the

emergence of this species and its multidrug resistance encourages new therapeutic strategies, including the combination of antifungals.

In summary, we demonstrated that there is no antagonism when flucytosine is combined *in vitro* with amphotericin B, voriconazole, or micafungin.

MATERIALS AND METHODS

A total of 15 *C. auris* isolates were included, of which 12 were clinical isolates collected from Indian patients during the years 2009 to 2011 (10) and 3 were the control and type strains from Korea (CBS 12372 and CBS 12373) from 2004 to 2006 (40) and Japan (CBS 10913) from 2009 (4). All strains have been previously identified by molecular techniques (4, 10, 40, 41).

The interactions of flucytosine with either amphotericin B, micafungin, or voriconazole were investigated by using a microdilution checkerboard method based on the EUCAST reference technique with 96-well microtiter plates (38). Drug dilutions were prepared to obtain four times the final concentration. Final concentrations ranged from 0.008 to 0.5 $\mu\text{g/ml}$ for flucytosine, 0.008 to 4 $\mu\text{g/ml}$ for amphotericin B, 0.004 to 2 $\mu\text{g/ml}$ for micafungin, and 0.008 to 4 $\mu\text{g/ml}$ for voriconazole. For two-dimensional microplate preparation, 50 μl of each concentration of flucytosine was added into wells 1 to 11 of each column and then 50 μl of the partner drug was added into wells A to H of each line. Line H and column 11 contained the partner drug and flucytosine alone, respectively. Column 12 served as the growth control containing only 50 μl of double-strength RPMI-2% glucose with dimethyl sulfoxide (DMSO). Plates were incubated at 37°C for 24 h and read spectrophotometrically. A growth inhibition endpoint of 50% was used both for the drugs tested alone and in combination. *Candida parapsilosis* ATCC 22019 and *Candida krusei* ATCC 6258 were used as quality controls. The experiments were performed in duplicate. Drug-free microplates were also incubated after inoculation with three strains, including the type strain, to ensure that possible heterogenous growth across the microplates (i.e., lower growth in the center area of the plate compared to outer wells) was not responsible for the observed effect (synergy or antagonism).

Two different methods were used to interpret the interactions of drugs. First, the fractional inhibitory concentration index (FICI) was calculated. The FICI was defined as $\text{FICI}_A + \text{FICI}_B = (C_A/\text{MIC}_A) + (C_B/\text{MIC}_B)$ where MIC_A and MIC_B are the MICs of drugs A and B alone and C_A and C_B are the concentrations of the drugs in combination, in all wells corresponding to an MIC. The interaction was considered synergistic when the FICI was ≤ 0.5 , indifferent at > 0.5 to ≤ 4.0 , and antagonistic at > 4 (42). An alternative approach, independent of an endpoint, was also used for visualization of the interaction between drugs. Briefly, data from combinations were processed for a response surface analysis by the Combenefit software, using the Bliss independence model (43, 44).

ACKNOWLEDGMENTS

During the past 5 years, E. Dannaoui has received research grants from MSD and Gilead; travel grants from Gilead, MSD, Pfizer, and Astellas; and speaker's fees from Gilead, MSD, and Astellas. F. Botterel has received research grants from MSD; travel grants from Gilead, MSD, and Pfizer; and speaker's fees from Gilead, MSD, and Pfizer.

REFERENCES

- Centers for Disease Control and Prevention. 2016. Clinical alert to U.S. healthcare facilities: global emergence of invasive infections caused by the multidrug-resistant yeast *Candida auris*. Centers for Disease Control and Prevention, Atlanta, GA. <https://www.cdc.gov/fungal/diseases/candidiasis/candida-auris-alert.html>.
- Spivak ES, Hanson KE. 2017. *Candida auris*: an emerging fungal pathogen. *J Clin Microbiol* 56:e01588-17. <https://doi.org/10.1128/JCM.01588-17>.
- Jeffery-Smith A, Taori SK, Schelenz S, Jeffery K, Johnson EM, Borman A, *Candida auris* Incident Management Team, Manuel R, Brown CS. 2017. *Candida auris*: a review of the literature. *Clin Microbiol Rev* 31:e00029-17. <https://doi.org/10.1128/CMR.00029-17>.
- Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. 2009. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol Immunol* 53:41–44. <https://doi.org/10.1111/j.1348-0421.2008.00083.x>.
- Lee WG, Shin JH, Uh Y, Kang MG, Kim SH, Park KH, Jang H-C. 2011. First three reported cases of nosocomial fungemia caused by *Candida auris*. *J Clin Microbiol* 49:3139–3142. <https://doi.org/10.1128/JCM.00319-11>.
- Ruiz Gaitán AC, Moret A, López Hontangas JL, Molina JM, Alexandre López AI, Cabezas AH, Mollar Maseres J, Arcas RC, Gómez Ruiz MD, Chiveli MÁ, Cantón E, Pemán J. 2017. Nosocomial fungemia by *Candida auris*: first four reported cases in continental Europe. *Rev Iberoam Micol* 34:23–27. <https://doi.org/10.1016/j.riam.2016.11.002>.
- Riat A, Neofytos D, Coste A, Harbarth S, Bizzini A, Grandbastien B, Pugin J, Lamoth F. 2018. First case of *Candida auris* in Switzerland: discussion about preventive strategies. *Swiss Med Wkly* 148:w14622. <https://doi.org/10.4414/SMW.2018.14622>.
- Larkin E, Hager C, Chandra J, Mukherjee PK, Retuerto M, Salem I, Long L, Isham N, Kovanda L, Borroto-Esoda K, Wring S, Angulo D, Ghannoum M. 2017. The emerging pathogen *Candida auris*: growth phenotype, virulence factors, activity of antifungals, and effect of SCY-078, a novel glucan synthesis inhibitor, on growth morphology and biofilm formation. *Antimicrob Agents Chemother* 61:e02396-16. <https://doi.org/10.1128/AAC.02396-16>.
- Bidaud AL, Chowdhary A, Dannaoui E. 2018. *Candida auris*: an emerging drug resistant yeast—a mini-review. *J Mycol Med* 28:568–573. <https://doi.org/10.1016/j.mycmed.2018.06.007>.
- Chowdhary A, Sharma C, Duggal S, Agarwal K, Prakash A, Singh PK, Jain S, Kathuria S, Randhawa HS, Hagen F, Meis JF. 2013. New clonal strain of *Candida auris*, Delhi, India. *Emerg Infect Dis* 19:1670–1673. <https://doi.org/10.3201/eid1910.130393>.
- Emara M, Ahmad S, Khan Z, Joseph L, Al-Obaid I, Purohit P, Bafna R. 2015. *Candida auris* candidemia in Kuwait, 2014. *Emerg Infect Dis* 21:1091–1092. <https://doi.org/10.3201/eid2106.150270>.
- Mohsin J, Hagen F, Al-Balushi ZAM, de Hoog GS, Chowdhary A, Meis JF, Al-Hatmi AMS. 2017. The first cases of *Candida auris* candidaemia in Oman. *Mycoses* 60:569–575. <https://doi.org/10.1111/myc.12647>.
- Araúz AB, Caceres DH, Santiago E, Armstrong P, Arosemena S, Ramos C, Espinosa-Bode A, Borace J, Hayer L, Cedeño I, Jackson BR, Sosa N,

- Berkow EL, Lockhart SR, Rodriguez-French A, Chiller T. 2018. Isolation of *Candida auris* from 9 patients in Central America: importance of accurate diagnosis and susceptibility testing. *Mycoses* 61:44–47. <https://doi.org/10.1111/myc.12709>.
14. Ben-Ami R, Berman J, Novikov A, Bash E, Shachor-Meyouhas Y, Zakin S, Maor Y, Tarabia J, Schechner V, Adler A, Finn T. 2017. Multidrug-resistant *Candida haemulonii* and *C. auris*, Tel Aviv, Israel. *Emerg Infect Dis* 23: 195–203. <https://doi.org/10.3201/eid2302.161486>.
 15. Chowdhary A, Anil Kumar V, Sharma C, Prakash A, Agarwal K, Babu R, Dinesh KR, Karim S, Singh SK, Hagen F, Meis JF. 2014. Multidrug-resistant endemic clonal strain of *Candida auris* in India. *Eur J Clin Microbiol Infect Dis* 33:919–926. <https://doi.org/10.1007/s10096-013-2027-1>.
 16. Centers for Disease Control and Prevention. 2019. Tracking *Candida auris*. Case count updated as of July 31, 2019. Centers for Disease Control and Prevention, Atlanta, GA. <https://www.cdc.gov/fungal/candida-auris/tracking-c-auris.html>.
 17. Wang X, Bing J, Zheng Q, Zhang F, Liu J, Yue H, Tao L, Du H, Wang Y, Wang H, Huang G. 2018. The first isolate of *Candida auris* in China: clinical and biological aspects. *Emerg Microbes Infect* 7:93. <https://doi.org/10.1038/s41426-018-0095-0>.
 18. Desoubeaux G, Bailly É, Guillaume C, De Kyvon M-A, Tellier A-C, Morange V, Bernard L, Salamé E, Quentin R, Chandenier J. 2018. *Candida auris* in contemporary mycology labs: a few practical tricks to identify it reliably according to one recent French experience. *J Mycol Med* 28:407–410. <https://doi.org/10.1016/j.mycmed.2018.02.011>.
 19. Mizusawa M, Miller H, Green R, Lee R, Durante M, Perkins R, Hewitt C, Simner PJ, Carroll KC, Hayden RT, Zhang SX. 2017. Can multidrug-resistant *Candida auris* be reliably identified in clinical microbiology laboratories? *J Clin Microbiol* 55:638–640. <https://doi.org/10.1128/JCM.02202-16>.
 20. Chowdhary A, Sharma C, Meis JF. 2017. *Candida auris*: a rapidly emerging cause of hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog* 13:e1006290. <https://doi.org/10.1371/journal.ppat.1006290>.
 21. Kathuria S, Singh PK, Sharma C, Prakash A, Masih A, Kumar A, Meis JF, Chowdhary A. 2015. Multidrug-resistant *Candida auris* misidentified as *Candida haemulonii*: characterization by matrix-assisted laser desorption ionization-time of flight mass spectrometry and DNA sequencing and its antifungal susceptibility profile variability by Vitek 2, CLSI broth microdilution, and Etest method. *J Clin Microbiol* 53:1823–1830. <https://doi.org/10.1128/JCM.00367-15>.
 22. Arendrup MC, Patterson TF. 2017. Multidrug-resistant *Candida*: epidemiology, molecular mechanisms, and treatment. *J Infect Dis* 216: S445–S451. <https://doi.org/10.1093/infdis/jix131>.
 23. Chowdhary A, Prakash A, Sharma C, Kordalewska M, Kumar A, Sarma S, Tarai B, Singh A, Upadhyaya G, Upadhyay S, Yadav P, Singh PK, Khillan V, Sachdeva N, Perlin DS, Meis JF. 2018. A multicentre study of antifungal susceptibility patterns among 350 *Candida auris* isolates (2009–17) in India: role of the ERG11 and FKS1 genes in azole and echinocandin resistance. *J Antimicrob Chemother* 73:891–899. <https://doi.org/10.1093/jac/dkx480>.
 24. Public Health England. 2017. Guidance for the laboratory investigation, management and infection prevention and control for cases of *Candida auris*. Public Health England, London, England. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/637685/Updated_Candida_auris_Guidance_v2.pdf.
 25. Centers for Disease Control and Prevention. 2019. Treatment and management of infections and colonization. Recommendations for treatment of *Candida auris* infections. Centers for Disease Control and Prevention, Atlanta, GA. <https://www.cdc.gov/fungal/candida-auris/c-auris-treatment.html>.
 26. Lockhart SR. 2019. *Candida auris* and multidrug resistance: defining the new normal. *Fungal Genet Biol* 131:103243. <https://doi.org/10.1016/j.fgb.2019.103243>.
 27. Schwarz P, Dromer F, Lortholary O, Dannaoui E. 2003. *In vitro* interaction of flucytosine with conventional and new antifungals against *Cryptococcus neoformans* clinical isolates. *Antimicrob Agents Chemother* 47: 3361–3364. <https://doi.org/10.1128/aac.47.10.3361-3364.2003>.
 28. Perfect JR, Dismukes WE, Dromer F, Goldman DL, Graybill JR, Hamill RJ, Harrison TS, Larsen RA, Lortholary O, Nguyen M-H, Pappas PG, Powderly WG, Singh N, Sobel JD, Sorrell TC. 2010. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis* 50:291–322. <https://doi.org/10.1086/649858>.
 29. Cornely OA, Bassetti M, Calandra T, Garbino J, Kullberg BJ, Lortholary O, Meersseman W, Akova M, Arendrup MC, Arkan-Akdagli S, Bille J, Castagnola E, Cuenca-Estrella M, Donnelly JP, Groll AH, Herbrecht R, Hope WW, Jensen HE, Lass-Flörl C, Petrikos G, Richardson MD, Roilides E, Verweij PE, Viscoli C, Ullmann AJ, ESCMID Fungal Infection Study Group. 2012. ESCMID guideline for the diagnosis and management of *Candida* diseases 2012: non-neutropenic adult patients. *Clin Microbiol Infect* 18:19–37. <https://doi.org/10.1111/1469-0691.12039>.
 30. Nguyen MH, Barchiesi F, McGough DA, Yu VL, Rinaldi MG. 1995. *In vitro* evaluation of combination of fluconazole and flucytosine against *Cryptococcus neoformans* var. *neoformans*. *Antimicrob Agents Chemother* 39:1691–1695. <https://doi.org/10.1128/aac.39.8.1691>.
 31. Mikami Y, Scalapone GM, Kurita N, Yazawa K, Uno J, Miyaji M. 1992. Synergistic postantifungal effect of flucytosine and fluconazole on *Candida albicans*. *J Med Vet Mycol* 30:197–206. <https://doi.org/10.1080/02681219280000261>.
 32. Polak A. 1987. Combination therapy of experimental candidiasis, cryptococcosis, aspergillosis and wangielliosis in mice. *Chemotherapy* 33: 381–395. <https://doi.org/10.1159/000238524>.
 33. Larsen RA, Bauer M, Weiner JM, Diamond DM, Leal ME, Ding JC, Rinaldi MG, Graybill JR. 1996. Effect of fluconazole on fungicidal activity of flucytosine in murine cryptococcal meningitis. *Antimicrob Agents Chemother* 40:2178–2182. <https://doi.org/10.1128/AAC.40.9.2178>.
 34. Fakhim H, Chowdhary A, Prakash A, Vaezi A, Dannaoui E, Meis JF, Badali H. 2017. *In vitro* interactions of echinocandins with triazoles against multidrug-resistant *Candida auris*. *Antimicrob Agents Chemother* 61: e01056-17. <https://doi.org/10.1128/AAC.01056-17>.
 35. de Oliveira HC, Monteiro MC, Rossi SA, Pemán J, Ruiz-Gaitán A, Mendes-Giannini MJS, Mellado E, Zaragoza O. 2019. Identification of off-patent compounds that present antifungal activity against the emerging fungal pathogen *Candida auris*. *Front Cell Infect Microbiol* 9:83. <https://doi.org/10.3389/fcimb.2019.00083>.
 36. Eldesouky HE, Li X, Abutaleb NS, Mohammad H, Seleem MN. 2018. Synergistic interactions of sulfamethoxazole and azole antifungal drugs against emerging multidrug-resistant *Candida auris*. *Int J Antimicrob Agents* 52:754–761. <https://doi.org/10.1016/j.ijantimicag.2018.08.016>.
 37. Lockhart SR, Etienne KA, Vallabhaneni S, Farooqi J, Chowdhary A, Govennder NP, Colombo AL, Calvo B, Cuomo CA, Desjardins CA, Berkow EL, Castanheira M, Magobo RE, Jabeen K, Asghar RJ, Meis JF, Jackson B, Chiller T, Litvintseva AP. 2017. Simultaneous emergence of multidrug-resistant *Candida auris* on 3 continents confirmed by whole-genome sequencing and epidemiological analyses. *Clin Infect Dis* 64:134–140. <https://doi.org/10.1093/cid/ciw691>.
 38. Arendrup MC, Prakash A, Meletiadis J, Sharma C, Chowdhary A. 2017. Comparison of EUCAST and CLSI reference microdilution MICs of eight antifungal compounds for *Candida auris* and associated tentative epidemiological cutoff values. *Antimicrob Agents Chemother* 61:e00485-17. <https://doi.org/10.1128/AAC.00485-17>.
 39. Chow NA, de Groot T, Badali H, Abastabar M, Chiller TM, Meis JF. 2019. Potential fifth clade of *Candida auris*, Iran, 2018. *Emerg Infect Dis* 25: 1780–1781. <https://doi.org/10.3201/eid2509.190686>.
 40. Kim M-N, Shin JH, Sung H, Lee K, Kim E-C, Ryoo N, Lee J-S, Jung S-I, Park KH, Kee SJ, Kim SH, Shin MG, Suh SP, Ryang DW. 2009. *Candida haemulonii* and closely related species at 5 university hospitals in Korea: identification, antifungal susceptibility, and clinical features. *Clin Infect Dis* 48:e57–e61. <https://doi.org/10.1086/597108>.
 41. Cendejas-Bueno E, Kolecka A, Alastruey-Izquierdo A, Theelen B, Groenewald M, Kostrzewa M, Cuenca-Estrella M, Gomez-Lopez A, Boekhout T. 2012. Reclassification of the *Candida haemulonii* complex as *Candida haemulonii* (C. *haemulonii* group I), *C. duobushaemulonii* sp. nov. (C. *haemulonii* group II), and *C. haemulonii* var. *vulnera* var. nov.: three multiresistant human pathogenic yeasts. *J Clin Microbiol* 50:3641–3651. <https://doi.org/10.1128/JCM.02248-12>.
 42. Odds FC. 2003. Synergy, antagonism, and what the checkerboard puts between them. *J Antimicrob Chemother* 52:1. <https://doi.org/10.1093/jac/dkg301>.
 43. Di Veroli GY, Fornari C, Wang D, Mollard S, Bramhall JL, Richards FM, Jodrell DI. 2016. Combeneft: an interactive platform for the analysis and visualization of drug combinations. *Bioinformatics* 32:2866–2868. <https://doi.org/10.1093/bioinformatics/btw230>.
 44. Schwarz P, Schwarz PV, Felske-Zech H, Dannaoui E. 2019. *In vitro* interactions between isavuconazole and tacrolimus, cyclosporin A or sirolimus against Mucorales. *J Antimicrob Chemother* 74:1921–1927. <https://doi.org/10.1093/jac/dkz102>.