



HAL
open science

Evaluation of the Gradient Concentration Strip Method for Antifungal Susceptibility Testing of Isavuconazole and Comparators for Mucorales Species

Pauline Vidal, Patrick Schwarz, Eric Dannaoui

► **To cite this version:**

Pauline Vidal, Patrick Schwarz, Eric Dannaoui. Evaluation of the Gradient Concentration Strip Method for Antifungal Susceptibility Testing of Isavuconazole and Comparators for Mucorales Species. *Antimicrobial Agents and Chemotherapy*, 2019, 63 (10), 10.1128/AAC.00838-19 . hal-03997639

HAL Id: hal-03997639

<https://hal.u-pec.fr/hal-03997639v1>

Submitted on 4 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Evaluation of the Gradient Concentration Strip Method for Antifungal Susceptibility Testing of Isavuconazole and Comparators for *Mucorales* Species

Pauline Vidal,^{a,b}  Patrick Schwarz,^{c,d}  Eric Dannaoui^{a,e,f}

^aUnité de Parasitologie-Mycologie, Service de Microbiologie, Hôpital Européen Georges Pompidou, Paris, France

^bFaculté de Pharmacie, Université Paris-Sud, Châtenay-Malabry, France

^cDepartment of Internal Medicine, Respiratory and Critical Care Medicine, University Hospital Marburg, Marburg, Germany

^dCenter for Invasive Mycoses and Antifungals, Philipps University Marburg, Marburg, Germany

^eFaculté de Médecine, Université Paris Descartes, Paris, France

^fDynamyc Research Group (EA 7380), Paris Est Créteil University, Ecole Nationale Vétérinaire d'Alfort, Créteil, France

ABSTRACT MIC values for amphotericin B and three azoles determined by the EUCAST reference technique and by gradient concentration strips were compared for 30 *Mucorales* isolates belonging to clinically important species. Essential agreement (EA) within ± 2 dilution steps at 24 hours between the techniques was 83.3% for isavuconazole. EAs for itraconazole, amphotericin B, and posaconazole were 86.7%, 73.3%, and 56.7%, respectively. A good agreement was obtained between visual and spectrophotometric readings for EUCAST.

KEYWORDS EUCAST, Etest, *Mucorales*, antifungal agents, antifungal susceptibility testing, filamentous fungi

Mucormycosis is associated with a high mortality rate, mostly affecting immunocompromised patients (1). *Mucorales* species show high *in vitro* MICs for several antifungals and are resistant to voriconazole (2–4), the first-line therapy for invasive aspergillosis. Recommended treatment for mucormycosis is high-dose liposomal amphotericin B combined with early surgery, if possible (5). Isavuconazole, a new broad-spectrum azole, is effective in patients with mucormycosis (6), generally well tolerated (7), and indicated for the treatment of mucormycosis when amphotericin B is not suitable (8). The recommended but sophisticated and time-consuming technique for antifungal susceptibility testing of molds is the broth microdilution method by CLSI or EUCAST (9, 10). Gradient concentration strips provide a fast and easy-to-handle tool for susceptibility testing of bacteria (11) and fungi (12). Whether antifungal MICs for *Mucorales* species obtained by the reference technique and the gradient concentration strips are comparable has been evaluated in a few studies with variable results (13–22). Data for isavuconazole are lacking.

(Part of this work was presented at the 28th European Congress of Clinical Microbiology and Infectious Diseases [ECCMID], 21 to 24 April 2018, Madrid, Spain.)

The aim of this study was to compare the MICs obtained by EUCAST and gradient concentration strip methodologies for isavuconazole, amphotericin B, posaconazole, and itraconazole for *Mucorales*.

Thirty *Mucorales* isolates, including 9 *Rhizopus arrhizus* (comprising 1 *R. arrhizus* var. *deleamar*), 3 *Rhizopus microsporus*, 5 *Lichtheimia corymbifera*, 5 *Lichtheimia ramosa*, 4 *Mucor circinelloides*, and 4 *Rhizomucor pusillus*, were tested. Isolates were identified by sequencing of the internal transcribed spacer regions. Sequences were previously

Citation Vidal P, Schwarz P, Dannaoui E. 2019. Evaluation of the gradient concentration strip method for antifungal susceptibility testing of isavuconazole and comparators for *Mucorales* species. *Antimicrob Agents Chemother* 63:e00838-19. <https://doi.org/10.1128/AAC.00838-19>.

Copyright © 2019 American Society for Microbiology. All Rights Reserved.

Address correspondence to Eric Dannaoui, eric.dannaoui@aphp.fr.

Received 19 April 2019

Returned for modification 15 June 2019

Accepted 19 July 2019

Accepted manuscript posted online 29 July 2019

Published 23 September 2019

deposited in GenBank (23–26). Quality control isolates included *Aspergillus fumigatus* ATCC 204305, *Candida krusei* ATCC 6258, and *Candida parapsilosis* ATCC 22019.

MICs were determined for isavuconazole, itraconazole, posaconazole, and amphotericin B by gradient concentration strip and EUCAST techniques (9). Isavuconazole (Basilea Pharmaceutical, Ltd., Basel, Switzerland), itraconazole (Sigma-Aldrich, Saint-Quentin Fallavier, France), posaconazole (MSD, Kenilworth, NJ), and amphotericin B (Sigma-Aldrich) were obtained as pure powders, and stock solutions were prepared in dimethyl sulfoxide.

MIC determination was performed in flat-bottomed 96-well microtiter plates using RPMI 1640 (with L-glutamine, with pH indicator, but without bicarbonate) (Sigma-Aldrich) (9). Final concentrations ranged from 0.015 to 8 $\mu\text{g/ml}$. Isolates were cultured for 5 to 7 days, and spore suspensions were prepared in water. Spore suspensions were counted with a hemocytometer and adjusted to 2 to 5×10^6 CFU/ml. Trays were inoculated with 100 μl of the 1/10-diluted spore suspension. Plates were incubated at 37°C (30°C for *M. circinelloides*) for 24 h, and MICs were determined visually and spectrophotometrically (complete inhibition).

For the gradient concentration strip method, RPMI agar plates were inoculated with the same spore suspension adjusted to 10^6 CFU/ml, and strips of isavuconazole (Liofilchem, Roseto degli Abruzzi, Italy) and the other drugs (Etest, bioMérieux, Marcy l'Etoile, France) were placed on the agar. MICs were determined after 24 h of incubation at 37°C (30°C for *M. circinelloides*). MICs were read where the inhibition ellipse intersected the MIC scale of the strip. Overgrowth by filaments bending into the ellipse was ignored.

Experiments were performed twice. Results of the two methods were analyzed by providing essential agreement (EA) values within ± 2 dilution steps. Agreement within ± 1 dilution step was also calculated. Categorical agreement was not calculated because there are currently no clinical breakpoints for *Mucorales* species.

Results of the *in vitro* susceptibility of *Mucorales* isolates determined by the EUCAST (visual reading) and gradient concentration strip methods are presented in Table 1. With the EUCAST method, isavuconazole MICs ranged from 0.25 to 16 $\mu\text{g/ml}$, with a geometric mean of 1.82 $\mu\text{g/ml}$. *In vitro* activity of isavuconazole depended on the species. The lowest and highest isavuconazole MICs were seen for *Lichtheimia* and *Mucor* isolates, respectively. With the gradient concentration strip method, isavuconazole MICs ranged from 0.25 to 16 $\mu\text{g/ml}$, with a geometric mean of 1.15 $\mu\text{g/ml}$. The EA between both techniques for isavuconazole was 83.3% (Table 2). When discrepant results were observed, gradient concentration strips showed lower MICs than the EUCAST method. Examples of results with the gradient concentration strips and the EUCAST inhibition curves are presented in Fig. 1.

For the comparators, MICs determined by the EUCAST method ranged from 0.125 to 2 $\mu\text{g/ml}$ (geometric mean, 0.26 $\mu\text{g/ml}$) for amphotericin B, 0.0625 to 16 $\mu\text{g/ml}$ (geometric mean, 1.66 $\mu\text{g/ml}$) for posaconazole, and 0.0625 to 16 $\mu\text{g/ml}$ (geometric mean, 1.7 $\mu\text{g/ml}$) for itraconazole. MICs determined by gradient concentration strips ranged from 0.125 to 16 $\mu\text{g/ml}$ (geometric mean, 0.81 $\mu\text{g/ml}$) for amphotericin B, 0.25 to 16 $\mu\text{g/ml}$ (geometric mean, 0.45 $\mu\text{g/ml}$) for posaconazole, and 0.03125 to 16 $\mu\text{g/ml}$ (geometric mean, 6.96 $\mu\text{g/ml}$) for itraconazole. EAs (± 2 dilution steps) for the comparators were 73.3%, 56.7%, and 86.7% for amphotericin B, posaconazole, and itraconazole, respectively (Table 2). Agreement within ± 1 dilution step was lower for isavuconazole, amphotericin B, posaconazole, and itraconazole at 66.7%, 53.3%, 40%, and 53.3%, respectively.

When MICs were determined spectrophotometrically with the EUCAST method, the EAs between the EUCAST and gradient concentration strip methods were 80%, 76.7%, 63.3%, and 73.3% for isavuconazole, amphotericin B, posaconazole, and itraconazole, respectively (data not shown).

The EAs between spectrophotometric and visual reading for the EUCAST technique were 93.3%, 100%, 86.2%, and 88.9% for isavuconazole, amphotericin B, posaconazole, and itraconazole, respectively (Table 3).

TABLE 1 Comparison of MICs determined visually by EUCAST broth susceptibility testing and by gradient concentration strip methodology for *Mucorales* isolates

Species	MIC ($\mu\text{g/ml}$) (range [geometric mean]) with ^a :									
	EUCAST method for:					Gradient concentration strip for:				
	ISA	AMB	PSZ	ITZ	AMB	ISA	AMB	PSZ	ITZ	AMB
<i>Rhizopus</i> spp. (n = 12)	1-8 (1.6)	0.25-2 (0.56)	0.125-2 (0.45)	4-16 (7)	0.25-4 (1.12)	0.5-16 (2.7)	0.25-16 (3.2)	0.5-16 (6.7)	0.25-2 (0.66)	0.25-16 (3.2)
<i>Lichtheimia</i> spp. (n = 10)	0.25-2 (1)	0.25-1 (0.15)	0.06-0.5 (0.134)	0.06-1 (0.33)	0.25-2 (0.6)	0.25-1 (0.66)	0.25-1 (0.57)	0.5-2 (0.9)	0.125	0.25-1 (0.57)
<i>Mucor circinelloides</i> (n = 4)	8-16 (11)	0.125-0.25 (0.17)	1-16 (4)	16	16	0.125	16	16	0.125	16
<i>Rhizomucor pusillus</i> (n = 4)	1-4 (1.6)	0.125-0.25 (0.15)	0.25-2 (0.5)	0.06-0.5 (0.15)	0.25-0.5 (0.42)	0.125-0.25 (0.25)	0.25-0.5 (0.35)	0.03-0.125 (0.07)	0.125-0.25 (0.25)	0.25-0.5 (0.35)
Total (n = 30)	0.25-16 (1.82)	0.125-2 (0.26)	0.06-16 (1.66)	0.06-16 (1.7)	0.25-16 (1.15)	0.125-16 (0.81)	0.25-16 (0.45)	0.03-16 (6.96)	0.125-16 (0.81)	0.25-16 (0.45)

^aISA, isavuconazole; AMB, amphotericin B; PSZ, posaconazole; ITZ, itraconazole.

TABLE 2 MIC dilution differences between EUCAST and gradient concentration strip methodologies for 30 *Mucorales* isolates

Drug ^a	Etest-EUCAST dilution difference							EA (%)	
	<-2	-2	-1	0	+1	+2	>2	±1 dilution step	±2 dilution steps
ISA	5	4	5	8	7	1	0	66.7	83.3
AMB	0	0	2	7	7	6	8	53.3	73.3
PSZ	1	0	1	7	4	5	12	40	56.7
ITZ	2	4	1	7	8	6	2	53.3	86.7

^aISA, isavuconazole; AMB, amphotericin B; PSZ, posaconazole; ITZ, itraconazole; EA, essential agreement.

First-line treatment for mucormycosis is liposomal amphotericin B (27); isavuconazole may be used when amphotericin B is inappropriate (8). Amphotericin B, isavuconazole, and posaconazole show good *in vitro* activity against *Mucorales* (2, 4, 26). Activity of itraconazole is limited to *Lichtheimia* spp., *Rhizomucor* spp., and some *Syncephalastrum racemosum* isolates (2, 4, 26). Here, the best *in vitro* activity, determined by the EUCAST technique, was seen for amphotericin B, with the lowest MICs for *Lichtheimia* spp. and *R. pusillus*. MICs for azole antifungals were generally higher and in accordance with previous studies (2, 4, 26).

Recommended techniques for antifungal susceptibility testing of molds are the reference techniques of CLSI and EUCAST. The gradient concentration strip method has shown good correlation with the reference techniques for yeasts and filamentous fungi (28). Nevertheless, only a few studies evaluated the correlation between gradient concentration strips and CLSI (14, 16–22) or EUCAST (13, 15) methodology for *Mucorales* species. One study compared isavuconazole results obtained by the CLSI and gradient concentration strip techniques (17), but correlation between the EUCAST and gradient concentration strip methods has never been evaluated. Here, the EA between the gradient concentration strip and EUCAST techniques for isavuconazole was 83.3%. When discrepant results were observed, except for *M. circinelloides*, gradient concentration strips gave lower MICs. The EAs between the two techniques were 73.3%, 56.7%, and 86.7% for amphotericin B, posaconazole, and itraconazole, respectively. Agreement




FIG 1 Examples of gradient concentration strip results and EUCAST inhibition curves obtained by spectrophotometric reading for an isolate of *L. corymbifera* (CBS 120581) for isavuconazole and comparators. Spectrophotometric curves were modeled by nonlinear regression based on a maximum-effect model with GraphPad Prism.

TABLE 3 Visual versus spectrophotometric reading of MICs determined by EUCAST methodology at 24 h

Drug ^a	Visual vs spectrophotometric MIC reading							EA (%)	
	<-2	-2	-1	0	+1	+2	>2	±1 dilution step	±2 dilution steps
ISA	0	1	1	13	11	2	2	83.3	93.3
AMB	0	1	0	19	9	1	0	93.3	100
PSZ	1	0	3	17	6	0	3	86.7	86.7
ITZ	0	0	0	18	6	2	4	80.0	86.7

^aISA, isavuconazole; AMB, amphotericin B; PSZ, posaconazole; ITZ, itraconazole; EA, essential agreement.

rates for the ± 1 dilution step were lower and in accordance with previous studies (13, 15, 17, 21, 22). Various results have been obtained for EAs between gradient concentration strip and CLSI or EUCAST techniques for *Mucorales* species (13–22). EAs ranged from 70.5% to 96.5% for amphotericin B (13–16, 19, 21, 22), 50% to 83% for itraconazole (18, 20–22), and 77% to 100% for posaconazole (13–16, 18, 19, 22). For isavuconazole, comparison of gradient concentration strips and the CLSI method resulted in an EA of 84.4% (17), similar to our results. Why the correlation between the techniques is sometimes not optimal remains unknown. Others have reported MIC reading difficulties for *Mucorales* isolates, because hyphae frequently overgrow the strip (13, 29). Incubation time may also be important, because better correlation was obtained for posaconazole with shorter incubation time (16), possibly related to different optimal growth temperatures of *Mucorales* species (30).

One of the drawbacks of the EUCAST technique for testing molds is the visual determination of MICs. Previous reports demonstrated that spectrophotometric reading is a good alternative for *Aspergillus* (31–33) and *Mucorales* (4) species. Here, correlation between visual determination and spectrophotometric reading for *Mucorales* isolates was high. Incorporation of spectrophotometric reading would lead to more objective MIC determination and automation of the process.

In conclusion, the EA between gradient concentration strips and EUCAST methodology for determination of isavuconazole MICs against *Mucorales* isolates was high (83.3%), although it did not reach the threshold of $\geq 90\%$. Further work is warranted to test the technical parameters that may improve the correlation between the two techniques.

ACKNOWLEDGMENTS

We are grateful to Basilea Pharmaceutica for providing the gradient concentration strips of isavuconazole.

P.V. has no conflicts of interest to declare. P.S. was supported by research grants from Basilea Pharmaceutica, Pfizer, and Gilead and received travel grants from Gilead and Pfizer. During the past 5 years, E.D. received research grants from MSD and Gilead, travel grants from Gilead, MSD, Pfizer, and Astellas, and speaker's fees from Gilead, MSD, and Astellas.

REFERENCES

- Petrikkos G, Skiada A, Drogari-Apiranthitou M. 2014. Epidemiology of mucormycosis in Europe. *Clin Microbiol Infect* 20(Suppl 6):67–73. <https://doi.org/10.1111/1469-0691.12563>.
- Alastruey-Izquierdo A, Castelli MV, Cuesta I, Zaragoza O, Monzon A, Mellado E, Rodriguez-Tudela JL. 2009. *In vitro* activity of antifungals against Zygomycetes. *Clin Microbiol Infect* 15(Suppl 5):71–76. <https://doi.org/10.1111/j.1469-0691.2009.02984.x>.
- Dannaoui E. 2017. Antifungal resistance in Mucorales. *Int J Antimicrob Agents* 50:617–621. <https://doi.org/10.1016/j.ijantimicag.2017.08.010>.
- Dannaoui E, Meletiadis J, Mouton JW, Meis JF, Verweij PE. 2003. *In vitro* susceptibilities of zygomycetes to conventional and new antifungals. *J Antimicrob Chemother* 51:45–52. <https://doi.org/10.1093/jac/dkg020>.
- Cornely OA, Arikan-Akdagli S, Dannaoui E, Groll AH, Lagrou K, Chakrabarti A, Lanternier F, Pagano L, Skiada A, Akova M, Arendrup MC, Boekhout T, Chowdhary A, Cuenca-Estrella M, Freiberger T, Guinea J, Guarro J, de Hoog S, Hope W, Johnson E, Kathuria S, Lackner M, Lass-Flörl C, Lortholary O, Meis JF, Meletiadis J, Munoz P, Richardson M, Roilides E, Tortorano AM, Ullmann AJ, van Diepeningen A, Verweij P, Petrikos G, European Society of Clinical Microbiology and Infectious Diseases Fungal Infection Study Group, European Confederation of Medical Mycology. 2014. ESCMID and ECMM joint clinical guidelines for the diagnosis and management of mucormycosis 2013. *Clin Microbiol Infect* 20(Suppl 3):5–26. <https://doi.org/10.1111/1469-0691.12371>.
- Marty FM, Ostrosky-Zeichner L, Cornely OA, Mullane KM, Perfect JR, Thompson GR, Alangaden GJ, Brown JM, Fredricks DN, Heinz WJ, Herbrecht R, Klimko N, Klyasova G, Maertens JA, Melinkeri SR, Oren I, Pappas PG, Ráčil Z, Rahav G, Santos R, Schwartz S, Vehreschild JJ, Young J-AH, Chetchotisakd P, Jaruratanasirikul S, Kanj SS, Engelhardt M, Kaufhold A,

- Ito M, Lee M, Sasse C, Maher RM, Zeiher B, Vehreschild MJGT, VITAL and FungiScope Mucormycosis Investigators. 2016. Isavuconazole treatment for mucormycosis: a single-arm open-label trial and case-control analysis. *Lancet Infect Dis* 16:828–837. [https://doi.org/10.1016/S1473-3099\(16\)00071-2](https://doi.org/10.1016/S1473-3099(16)00071-2).
7. Natesan SK, Chandrasekar PH. 2016. Isavuconazole for the treatment of invasive aspergillosis and mucormycosis: current evidence, safety, efficacy, and clinical recommendations. *Infect Drug Resist* 9:291–300. <https://doi.org/10.2147/IDR.S102207>.
 8. Sipsas NV, Gamaletsou MN, Anastasopoulou A, Kontoyiannis DP. 2018. Therapy of mucormycosis. *J Fungi (Basel)* 4:E90. <https://doi.org/10.3390/jof4030090>.
 9. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope W, Howard SJ. 2014. Method for the determination of broth dilution minimum inhibitory concentrations of antifungal agents for conidia forming moulds. EUCAST definitive document EDef 9.2. EUCAST, Basel, Switzerland.
 10. Clinical and Laboratory Standards Institute. 2008. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi. CLSI document M38-A2. Clinical and Laboratory Standards Institute, Wayne, PA.
 11. Jorgensen JH, Ferraro MJ. 2009. Antimicrobial susceptibility testing: a review of general principles and contemporary practices. *Clin Infect Dis* 49:1749–1755. <https://doi.org/10.1086/647952>.
 12. Lass-Flörl C, Perkhof S, Mayr A. 2010. In vitro susceptibility testing in fungi: a global perspective on a variety of methods. *Mycoses* 53:1–11. <https://doi.org/10.1111/j.1439-0507.2009.01813.x>.
 13. Caramalho R, Maurer E, Binder U, Araújo R, Dolatabadi S, Lass-Flörl C, Lackner M. 2015. Etest cannot be recommended for *in vitro* susceptibility testing of *Mucorales*. *Antimicrob Agents Chemother* 59:3663–3665. <https://doi.org/10.1128/AAC.00004-15>.
 14. Chowdhary A, Kathuria S, Singh PK, Sharma B, Dolatabadi S, Hagen F, Meis JF. 2014. Molecular characterization and *in vitro* antifungal susceptibility of 80 clinical isolates of mucormycetes in Delhi, India. *Mycoses* 57(Suppl 3):97–107. <https://doi.org/10.1111/myc.12234>.
 15. Drogari-Apiranthitou M, Mantopoulou FD, Skiada A, Kanioura L, Grammatikou M, Vrioni G, Mitroussia-Ziouva A, Tsakris A, Petrikkos G. 2012. In vitro antifungal susceptibility of filamentous fungi causing rare infections: synergy testing of amphotericin B, posaconazole and anidulafungin in pairs. *J Antimicrob Chemother* 67:1937–1940. <https://doi.org/10.1093/jac/dks137>.
 16. Espinel-Ingroff A. 2006. Comparison of three commercial assays and a modified disk diffusion assay with two broth microdilution reference assays for testing zygomycetes, *Aspergillus* spp., *Candida* spp., and *Cryptococcus neoformans* with posaconazole and amphotericin B. *J Clin Microbiol* 44:3616–3622. <https://doi.org/10.1128/JCM.01187-06>.
 17. Guinea J, Pelaez T, Recio S, Torres-Narbona M, Bouza E. 2008. *In vitro* antifungal activities of isavuconazole (BAL4815), voriconazole, and fluconazole against 1,007 isolates of zygomycete, *Candida*, *Aspergillus*, *Fusarium*, and *Scedosporium* species. *Antimicrob Agents Chemother* 52:1396–1400. <https://doi.org/10.1128/AAC.01512-07>.
 18. Kondori N, Svensson E, Mattsby-Baltzer I. 2011. In vitro susceptibility of filamentous fungi to itraconazole, voriconazole and posaconazole by Clinical and Laboratory Standards Institute reference method and E-test. *Mycoses* 54:e318. <https://doi.org/10.1111/j.1439-0507.2010.01913.x>.
 19. Lamoth F, Alexander BD. 2015. Comparing Etest and broth microdilution for antifungal susceptibility testing of the most-relevant pathogenic molds. *J Clin Microbiol* 53:3176–3181. <https://doi.org/10.1128/JCM.00925-15>.
 20. Pfaller MA, Messer SA, Mills K, Bolmstrom A. 2000. *In vitro* susceptibility testing of filamentous fungi: comparison of Etest and reference microdilution methods for determining itraconazole MICs. *J Clin Microbiol* 38:3359–3361.
 21. Szekely A, Johnson EM, Warnock DW. 1999. Comparison of E-test and broth microdilution methods for antifungal drug susceptibility testing of molds. *J Clin Microbiol* 37:1480–1483.
 22. Torres-Narbona M, Guinea J, Martínez-Alarcón J, Peláez T, Bouza E. 2007. *In vitro* activities of amphotericin B, caspofungin, itraconazole, posaconazole, and voriconazole against 45 clinical isolates of zygomycetes: comparison of CLSI M38-A, Sensititre YeastOne, and the Etest. *Antimicrob Agents Chemother* 51:1126–1129. <https://doi.org/10.1128/AAC.01539-06>.
 23. Alastruey-Izquierdo A, Hoffmann K, de Hoog GS, Rodriguez-Tudela JL, Voigt K, Bibashi E, Walther G. 2010. Species recognition and clinical relevance of the zygomycetous genus *Lichtheimia* (syn. *Absidia pro parte*, *Mycocladius*). *J Clin Microbiol* 48:2154–2170. <https://doi.org/10.1128/JCM.01744-09>.
 24. Garcia-Hermoso D, Hoinard D, Gantier JC, Grenouillet F, Dromer F, Dannaoui E. 2009. Molecular and phenotypic evaluation of *Lichtheimia corymbifera* (formerly *Absidia corymbifera*) complex isolates associated with human mucormycosis: rehabilitation of *L. ramosa*. *J Clin Microbiol* 47:3862–3870. <https://doi.org/10.1128/JCM.02094-08>.
 25. Schwarz P, Bretagne S, Gantier JC, Garcia-Hermoso D, Lortholary O, Dromer F, Dannaoui E. 2006. Molecular identification of zygomycetes from culture and experimentally infected tissues. *J Clin Microbiol* 44:340–349. <https://doi.org/10.1128/JCM.44.2.340-349.2006>.
 26. Vitale RG, de Hoog GS, Schwarz P, Dannaoui E, Deng S, Machouart M, Voigt K, van de Sande WW, Dolatabadi S, Meis JF, Walther G. 2012. Antifungal susceptibility and phylogeny of opportunistic members of the order *Mucorales*. *J Clin Microbiol* 50:66–75. <https://doi.org/10.1128/JCM.06133-11>.
 27. Skiada A, Pagano L, Groll A, Zimmerli S, Dupont B, Lagrou K, Lass-Flörl C, Bouza E, Klimko N, Gaustad P, Richardson M, Hamal P, Akova M, Meis JF, Rodriguez-Tudela JL, Roilides E, Mitroussia-Ziouva A, Petrikkos G, European Confederation of Medical Mycology Working Group on Zygomycosis. 2011. Zygomycosis in Europe: analysis of 230 cases accrued by the registry of the European Confederation of Medical Mycology (ECMM) working group on zygomycosis between 2005 and 2007. *Clin Microbiol Infect* 17:1859–1867. <https://doi.org/10.1111/j.1469-0691.2010.03456.x>.
 28. Moser SA, Wicker J. 2016. Commercial methods for identification and susceptibility testing of fungi, p 214–272. *In* Truant AL (ed), *Manual of commercial methods in clinical microbiology*. John Wiley & Sons, Hoboken, NJ.
 29. Maurer E, Binder U, Sparber M, Lackner M, Caramalho R, Lass-Flörl C. 2015. Susceptibility profiles of amphotericin B and posaconazole against clinically relevant *Mucorales* species under hypoxic conditions. *Antimicrob Agents Chemother* 59:1344–1346. <https://doi.org/10.1128/AAC.04424-14>.
 30. de Hoog GS, Guarro J (ed). 1995. *Atlas of clinical fungi*. Centraalbureau voor Schimmelcultures, Baarn, The Netherlands.
 31. Dannaoui E, Lortholary O, Dromer F. 2004. *In vitro* evaluation of double and triple combinations of antifungal drugs against *Aspergillus fumigatus* and *Aspergillus terreus*. *Antimicrob Agents Chemother* 48:970–978. <https://doi.org/10.1128/aac.48.3.970-978.2004>.
 32. Dannaoui E, Persat F, Monier MF, Borel E, Piens MA, Picot S. 1999. Use of spectrophotometric reading for *in vitro* antifungal susceptibility testing of *Aspergillus* spp. *Can J Microbiol* 45:871–874. <https://doi.org/10.1139/w99-075>.
 33. Meletiadis J, Leth Mortensen K, Verweij PE, Mouton JW, Arendrup MC. 2017. Spectrophotometric reading of EUCAST antifungal susceptibility testing of *Aspergillus fumigatus*. *Clin Microbiol Infect* 23:98–103. <https://doi.org/10.1016/j.cmi.2016.10.017>.