



**HAL**  
open science

## Investigation of an Outbreak of COVID-19 in a French Nursing Home With Most Residents Vaccinated

Catherine Burugorri-Pierre, Carmelo Lafuente-Lafuente, Christel Oasi, Emmanuel Lecorche, Sylvie Pariel, Cristiano Donadio, Joël Belmin

► **To cite this version:**

Catherine Burugorri-Pierre, Carmelo Lafuente-Lafuente, Christel Oasi, Emmanuel Lecorche, Sylvie Pariel, et al.. Investigation of an Outbreak of COVID-19 in a French Nursing Home With Most Residents Vaccinated. *JAMA Network Open*, 2021, 4 (9), pp.e2125294. 10.1001/jamanetworkopen.2021.25294 . hal-04146198

**HAL Id: hal-04146198**

**<https://hal.u-pec.fr/hal-04146198v1>**

Submitted on 29 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Investigation of an Outbreak of COVID-19 in a French Nursing Home With Most Residents Vaccinated

Catherine Burugorri-Pierre, MD; Carmelo Lafuente-Lafuente, MD, PhD; Christel Oasi, MD; Emmanuel Lecorche, PharmD, PhD; Sylvie Pariel, MD; Cristiano Donadio, MD; Joël Belmin, MD, PhD

## Introduction

There is great hope that vaccination against SARS-CoV-2 will decrease the burden of COVID-19 on nursing home (NH) residents, who have been significantly affected by this pandemic. The efficacy and effectiveness of COVID-19 vaccines are not well known in this population. Recent immunogenic studies have found decreased titers of postvaccine neutralizing antibodies against SARS-CoV-2 among NH residents, suggesting the potential for diminished effectiveness in this population.<sup>1,2</sup> We report an outbreak of COVID-19 in a French nursing home where most residents had been fully vaccinated with the BNT162b2 vaccine.

## Methods

This cohort study was conducted according to the principle of the Declaration of Helsinki. The president of the Comité de Protection des Personnes Ile-de-France VI determined that this study was exempt from ethics committee review and oral or written participant consent according to the French law (décret No. 2016-1537, November 17, 2016). Residents of the facility and their families were informed that a report about the outbreak would be published using deidentified data. This report follows the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline for cohort studies.

The study reports an outbreak that occurred in a 77-bed NH located in Biscarrosse, France, in which a vaccine campaign was conducted using the BNT162b2 vaccine in January and February 2021. Several weeks after this campaign, a resident contracted COVID-19 outside of the facility. To control the spread of the infection, all residents and health care professionals who accepted the procedure repeatedly underwent nasal swabbing for reverse transcriptase-polymerase chain reaction (RT-PCR) testing for SARS-CoV-2. An outbreak of COVID-19 ensued, nonetheless. We followed up all individuals diagnosed with COVID-19 and extracted clinical data from residents' medical records. Information about health care professionals was obtained from administrative records kept by the NH director. Additional methods are detailed in eMethods in the [Supplement](#). These patients and data have not been previously reported. Data were analyzed from March 19 through April 18, 2021.

## Results

The index infection occurred in a resident who came into contact with a visitor who had symptoms. The visitor was diagnosed with COVID-19 and informed the facility of this diagnosis 4 days later. The resident was at that time asymptomatic with a positive RT-PCR test result, and she was isolated. She became symptomatic 2 days later and was treated with nasal oxygenation, fluid infusion, and anticoagulants.

The results of RT-PCR tests are presented in the [Table](#). Among 74 residents (16 men [22.2%]; mean [SD] age, 87.8 [7.5] years), 72 individuals (97.3%) were vaccinated, including 70 residents who were fully vaccinated (ie, received 2 doses of BNT162b2 >14 days before the outbreak) and 2

## + Supplemental content

Author affiliations and article information are listed at the end of this article.

residents who were partially vaccinated (ie, received 1 dose >14 days before the outbreak). In total, 17 residents (23.0%) were diagnosed with COVID-19 (5 men [29.4%] and 12 women [70.6%]; mean [SD] age, 87.6 [8.9] years). The individuals diagnosed with COVID-19 included 1 unvaccinated resident, 2 partially vaccinated residents, and 14 fully vaccinated residents. Among individuals with diagnoses, 8 residents developed severe disease, 2 were hospitalized, and 1 individual (the unvaccinated resident) died (Figure).

Among 102 health care professionals, 34 individuals were vaccinated (33.3%), 68 individuals were unvaccinated (66.7%), and 12 individuals were diagnosed with COVID-19 (11.8%). Of these, 5 individuals became symptomatic and none had severe disease. Among health care professionals who were unvaccinated, 9 individuals (13.2%) were diagnosed with COVID-19, and among health care professionals who were previously vaccinated, 3 individuals (8.8%) were diagnosed (including 1 individual fully vaccinated with the BNT162b2 vaccine and 2 individuals partially vaccinated with the ChAdOx1 nCoV-19 vaccine).

The SARS-CoV-2 variant B.1.1.7 was identified among all residents and health care professionals who were diagnosed with COVID-19. Viral genomes were sequenced for 9 samples obtained from 4 residents and 5 health care professionals. The profile of this variant revealed no specific mutations in the spike gene.


Table. RT-PCR Test Results, Confirmed Diagnoses, and Deaths by Vaccination Status

Outcome	Residents (n = 74)		Health care professionals (n = 102)	
	Vaccinated (n = 72)	Unvaccinated (n = 2)	Vaccinated (n = 34)	Unvaccinated (n = 68)
RT-PCR positive test result, No./No. total tests <sup>a</sup>				
Day 3	3/70	1/2	0/32	0/65
Day 7 or 8	10/59	0/1	1/34	4/68
Day 15	2/52	0/1	2/32	6/65
Confirmed COVID-19 diagnosis, No. (%)				
Any	16 (22.2)	1 (50.0)	3 (8.8)	9 (13.2)
Symptomatic COVID-19, No. (%)	14 (87.5)	1 (100)	0	5 (55.6)
Severe COVID-19, No. (%)	7 (50.0)	1 (100)	0	0
COVID-19-related death, No. (%)	0	1 (100)	0	0

Abbreviation: RT-PCR, reverse transcriptase-polymerase chain reaction.

<sup>a</sup> RT-PCR results are for tests performed after the index diagnosis (ie, day 0). Positive results are out of the total number of individuals receiving RT-PCR tests that day.

Figure. Timeline of Test Results, Symptoms, Symptom Duration, and Complications



Outcomes are among 17 nursing home residents with a confirmed diagnosis of COVID-19. Individual No. 4 was unvaccinated, and all others were vaccinated with BNT162b2 vaccine; individuals No. 7 and 17 had only 1 dose. RT-PCR indicates reverse transcriptase-polymerase chain reaction.

## Discussion

This cohort study's findings suggest that an outbreak of COVID-19 can occur among fully vaccinated NH residents. The study found evidence of transmission among vaccinated residents, but few individuals who were infected developed severe disease and 1 patient, who was unvaccinated, died. Moreover, these outcomes occurred in a setting in which approximately 30% of staff members were vaccinated.

To our knowledge, this is the first outbreak due to the B.1.1.7 COVID-19 variant described among individuals well vaccinated against SARS-CoV-2. Cavanaugh et al<sup>3</sup> reported a COVID-19 outbreak associated with a SARS-CoV-2 R.1 lineage variant that occurred in a US skilled nursing facility where 90% of residents were fully vaccinated.

The occurrence of the outbreak described in our study despite vaccination may be associated with impaired immune function associated with immunosenescence and health conditions, such as malnutrition, diabetes, and cancer, that are frequent among NH residents.<sup>4</sup> Several studies<sup>1,2,4-6</sup> have found reduced antibody response to several vaccines, including BNT162b2, in this population.

Our study's findings suggest that SARS-CoV-2 vaccination may not be sufficient as the sole means to prevent COVID-19 among NH residents and that other prevention measures should not be abandoned yet in these settings. More research is needed to improve the effectiveness of SARS-CoV-2 vaccines in this population.

---

### ARTICLE INFORMATION

**Accepted for Publication:** July 12, 2021.

**Published:** September 13, 2021. doi:10.1001/jamanetworkopen.2021.25294

**Open Access:** This is an open access article distributed under the terms of the [CC-BY License](#). © 2021 Burugorri-Pierre C et al. *JAMA Network Open*.

**Corresponding Author:** Joël Belmin, MD, PhD, Service de Gériatrie, Hôpital Charles Foix, Assistance Publique-Hopitaux de Paris, 7 avenue de la République, 94200 Ivry-sur-Seine, France ([j.belmin@aphp.fr](mailto:j.belmin@aphp.fr)).

**Author Affiliations:** Etablissement d'Hébergement pour Personnes Agées Dépendantes Léon Dubédac, Biscarrosse, France (Burugorri-Pierre); Service de Gériatrie, Hôpital Charles Foix, Assistance Publique-Hopitaux de Paris, Ivry-sur-Seine, France (Lafuente-Lafuente, Oasi, Pariel, Donadio, Belmin); Faculté de Médecine Sorbonne, Sorbonne Université, Paris, France (Lafuente-Lafuente, Belmin); Clinical Epidemiology and Ageing Unit, Institut National de la Santé et de la Recherche Médicale U955, Institut Mondor de Recherche Biomédicale, Créteil, France (Lafuente-Lafuente); Laboratoire Cerba, Saint-Ouen L'Aumône, France (Lecorche).

**Author Contributions:** Dr Burugorri-Pierre had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

*Concept and design:* Lafuente-Lafuente, Oasi, Donadio, Belmin.

*Acquisition, analysis, or interpretation of data:* Burugorri-Pierre, Lafuente-Lafuente, Lecorche, Pariel, Belmin.

*Drafting of the manuscript:* Burugorri-Pierre, Belmin.

*Critical revision of the manuscript for important intellectual content:* Lafuente-Lafuente, Oasi, Lecorche, Pariel, Donadio, Belmin.

*Statistical analysis:* Burugorri-Pierre, Lafuente-Lafuente, Lecorche.

*Administrative, technical, or material support:* Belmin.

*Supervision:* Lafuente-Lafuente, Belmin.

**Conflict of Interest Disclosures:** Dr Belmin reported receiving personal fees from Novartis, Pfizer, and Sanofi outside the submitted work. No other disclosures were reported.

**Additional Contributions:** Bénédicte Roquebert, PhD (Cerba Laboratory), contributed to viral genome sequencing, and Laurence Jacquenod, MA (Sorbonne Université), provided technical help with the Figure. These individuals were not compensated for this work.

**REFERENCES**

1. Van Praet JT, Vandecasteele S, De Roo A, De Vriese AS, Reynders M. Humoral and cellular immunogenicity of the BNT162b2 mRNA COVID-19 vaccine in nursing home residents. *Clin Infect Dis*. Published online ahead of print April 7, 2021;ciab300. doi:[10.1093/cid/ciab300](https://doi.org/10.1093/cid/ciab300)
2. Brockman MA, Mwimanzi F, Sang Y, et al. Weak humoral immune reactivity among residents of long-term care facilities following one dose of the BNT162b2 mRNA COVID-19 vaccine. *medRxiv*. Preprint posted online March 24, 2021:2021.03.17.21253773. doi:[10.1101/2021.03.17.21253773](https://doi.org/10.1101/2021.03.17.21253773)
3. Cavanaugh AM, Fortier S, Lewis P, et al. COVID-19 outbreak associated with a SARS-CoV-2 R.1 lineage variant in a skilled nursing facility after vaccination program—Kentucky, March 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(17):639-643. doi:[10.15585/mmwr.mm7017e2](https://doi.org/10.15585/mmwr.mm7017e2)
4. Fulop T, Pawelec G, Castle S, Loeb M. Immunosenescence and vaccination in nursing home residents. *Clin Infect Dis*. 2009;48(4):443-448. doi:[10.1086/596475](https://doi.org/10.1086/596475)
5. Lelic A, Verschoor CP, Lau VW, et al. Immunogenicity of varicella vaccine and immunologic predictors of response in a cohort of elderly nursing home residents. *J Infect Dis*. 2016;214(12):1905-1910. doi:[10.1093/infdis/jiw462](https://doi.org/10.1093/infdis/jiw462)
6. Williams RE, Sena AC, Moorman AC, et al. Hepatitis B vaccination of susceptible elderly residents of long term care facilities during a hepatitis B outbreak. *Vaccine*. 2012;30(21):3147-3150. doi:[10.1016/j.vaccine.2012.02.078](https://doi.org/10.1016/j.vaccine.2012.02.078)

**SUPPLEMENT.****eMethods.**