

A multi-scale model for CD8+ T cell immune response

Fabien Crauste, Olivier Gandrillon, Madge Martin, Thi Nhu Thao Nguyen,

Samuel Bernard

▶ To cite this version:

Fabien Crauste, Olivier Gandrillon, Madge Martin, Thi Nhu Thao Nguyen, Samuel Bernard. A multi-scale model for CD8+ T cell immune response. Journées Math Bio Santé 2022, GDR MathSAV; GT Mabiome, Oct 2022, Besançon, France. hal-04193666

HAL Id: hal-04193666 https://hal.u-pec.fr/hal-04193666v1

Submitted on 1 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

A multi-scale model for CD8+ T cell immune response

Fabien Crauste¹, Olivier Gandrillon^{2,3}, Madge Martin⁴, Thi Nhu Thao Nguyen^{*2,3}, and Samuel Bernard^{3,5}

¹Laboratoire MAP5 – Université Paris Cité – Paris, France

²Laboratoire de biologie et modélisation de la cellule – École Normale Supérieure - Lyon – Lyon, France ³Equipe Inria Dracula – INRIA Lyon – Villeurbanne, France

⁴CNRS, Univ Paris Est Creteil, Univ Gustave Eiffel, UMR 8208 – Laboratoire MSME (Modélisation et Simulation Multi Echelle) – F-94010 Créteil, France

⁵Institut Camille Jordan [Villeurbanne] – Université Claude Bernard Lyon 1 – France

Résumé

CD8+ T cells are important for immune responses against viruses and bacteria, as well as for tumor surveillance. When the CD8+ T cell recognizes its antigen (i.e. it enters in contact with an antigen presenting cell), it gets activated, it then differentiates into an effector, and kills infected or malignant cells bearing the antigen. Meanwhile, part of activated CD8+ T cells differentiate into memory cells. CD8+ T cell activation, differentiation, proliferation and death are controlled by intracellular signaling based on cell interactions with their microenvironment and gene regulation (1), (2). Here we present a multiscale model for CD8+ T cell immune response.

The model is implemented in the modelling platform Simuscale developed at Inria. Simuscale provides an efficient numerical framework for integrating and simulating different physical and temporal scales involved in the CD8 T cell immune response. First, a gene regulatory network (GRN), described by a Piecewise-deterministic Markov process (PDMP), is integrated and simulated in each individual T cell. Second, the internal GRN dynamics affects cellular fate: it changes cell activation status and triggers either proliferation, apoptosis, or differentiation. Third, interactions between cells modify the GRN dynamics via cell signaling, generating feedback regulation of CD8 T cell dynamics.

Through simulations, we explore conditions (for example threshold values of the protein concentration of a given gene, or the volume of a cell) that can lead to decisions about division, death, and differentiation states of a cell. We introduce a GRN consisting of 8 different genes. Simulations of this complex, multiscale model show that the model is able to reproduce key features of the CD8 T cell response (3).

References

(1) Girel, S., et al., Model-based assessment of the role of uneven partitioning of molecular content on heterogeneity and regulation of differentiation in CD8 T-cell immune responses,

^{*}Intervenant

Front. Immunol, 2019.

(2) Gao, X., et al., IL-2 sensitivity and exogenous IL-2 concentration gradient tune the productive contact duration of CD8+ T cell-APC : a multiscale modeling study, BMC Sys Biol, 2016.

(3) Todorov, H., et al., CD8 memory precursor cells generation is a continuous process, iScience, 2022.